

Inflammatory Bowel Diseases

Interactions between Autophagy and the Unfolded Protein Response: Implications for Inflammatory Bowel Disease

--Manuscript Draft--

Manuscript Number:	IBD-D-18-00574R1
Article Type:	Review Article - Basic Science
Keywords:	Inflammatory bowel disease; Crohn's disease; autophagy; Unfolded Protein Response; ER Stress
Corresponding Author:	Craig Stevens, PhD Edinburgh Napier University Faculty of Health Life and Social Sciences Edinburgh, United Kingdom
First Author:	Craig Stevens, PhD
Order of Authors:	Craig Stevens, PhD Kirsty M. Hooper Peter G. Barlow Paul Henderson
Manuscript Region of Origin:	UNITED KINGDOM
Abstract:	<p>Inflammatory Bowel Disease (IBD), which includes Crohn's disease (CD) and ulcerative colitis, is characterised by chronic inflammation of the gastrointestinal tract. Aetiology involves a combination of genetic and environmental factors resulting in abnormal immune responses to intestinal microbiota. Genetic studies have strongly linked genes involved in autophagy to CD, and genes involved in the unfolded protein response (UPR) to IBD. The UPR is triggered in response to accumulation of misfolded proteins in the endoplasmic reticulum (ER) and autophagy plays a key role to relieve ER-stress and restore homeostasis. This review summarises the known interactions between autophagy and the UPR and discusses the impact of these converging pathways on IBD pathogenesis. With a paucity of effective long-term treatments for IBD, targeting of synergistic pathways may provide novel and more effective therapeutic options.</p>

1 1 Interactions between Autophagy and the Unfolded Protein

2 2 Response: Implications for Inflammatory Bowel Disease

3
4
5 3
6
7
8
9
10
11
12
13 4 Kirsty M. Hooper¹, Peter G. Barlow¹, Paul Henderson^{2, 3¶} and Craig Stevens^{1¶*}.

14
15
16 5
17
18 6 1. School of Applied Sciences, Edinburgh Napier University, Sighthill Campus, Sighthill Court,
19 7 Edinburgh, EH11 4BN.

20
21
22 8 2. Child Life and Health, University of Edinburgh, Edinburgh, EH9 1UW.

23
24
25 9 3. Department of Paediatric Gastroenterology and Nutrition, Royal Hospital for Sick Children,
26 10 Edinburgh, EH9 1LF.

27
28 11 ¶Joint senior authors

29
30
31 12
32
33
34 13 Short title: Autophagy and UPR in IBD

35
36 14
37
38 15 *Address for Correspondence:

39
40
41 16 Dr Craig Stevens

42
43
44 17 School of Applied Sciences, Edinburgh Napier University, Sighthill Campus, Sighthill Court,
45 18 Edinburgh, EH11 4BN.

46
47 19 Email: C.Stevens@napier.ac.uk

48
49
50 20 Tel: 0044 131 455 2930

51
52
53 21

54
55
56
57 22

58
59
60
61
62
63
64
65

23 Abstract

1
2
3
4
5 24 Inflammatory Bowel Disease (IBD), which includes Crohn's disease (CD) and ulcerative colitis,
6
7 25 is characterised by chronic inflammation of the gastrointestinal tract. Aetiology involves a
8
9
10 26 combination of genetic and environmental factors resulting in abnormal immune responses
11
12
13 27 to intestinal microbiota. Genetic studies have strongly linked genes involved in autophagy to
14
15 28 CD, and genes involved in the unfolded protein response (UPR) to IBD. The UPR is triggered
16
17
18 29 in response to accumulation of misfolded proteins in the endoplasmic reticulum (ER) and
19
20
21 30 autophagy plays a key role to relieve ER-stress and restore homeostasis. This review
22
23 31 summarises the known interactions between autophagy and the UPR and discusses the
24
25
26 32 impact of these converging pathways on IBD pathogenesis. With a paucity of effective long-
27
28
29 33 term treatments for IBD, targeting of synergistic pathways may provide novel and more
30
31 34 effective therapeutic options.

32
33
34
35 35 **Keywords:** IBD, autophagy, unfolded protein response, ER stress.
36
37
38
39 36
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

37 Introduction

38 Inflammatory Bowel Disease (IBD) is a group of inflammatory diseases that includes Crohn's
39 disease (CD), ulcerative colitis (UC) and IBD unclassified (IBDU). The incidence rate for IBD is
40 approximately 50-200 in 100,000 persons per year in Western countries [1] and following
41 diagnosis the natural history of the condition is characterized by periods of relapse and
42 remission, with symptoms commonly including abdominal pain, chronic diarrhoea, weight
43 loss and lethargy [2]. CD is distinguished from UC due to the presence of submucosal or
44 transmural inflammation and macroscopic changes that often occur in a non-contiguous
45 pattern anywhere within the digestive tract [1]. UC is localised to the colon and inflammation
46 is limited to the mucosa and epithelial lining of the gastrointestinal (GI) tract [2]. Patients can
47 be diagnosed with IBDU when a conclusive distinction between CD and UC cannot be made,
48 although this may well represent a distinct sub-type. At present there is no cure for IBD and
49 medications such as corticosteroids, aminosalicylates, immunomodulators and biological
50 agents are aimed at inducing and maintaining remission of disease by modifying inflammatory
51 processes [3].

52 The aetiopathology of IBD is multifactorial in nature, with genetic predisposition,
53 environmental triggers (e.g. smoking, appendectomy, diet, pollution, antibiotics and stress)
54 and a dysregulated mucosal immune response contributing to disease [4]. Examination of the
55 gut microbiome has revealed that IBD is associated with microbial dysbiosis, including an
56 expansion of facultative anaerobic bacteria of the family Enterobacteriaceae [5]. Several
57 potentially causative agents have been identified, most notably *Escherichia coli* strains with
58 an adherent and invasive phenotype (AIEC) are associated with ileal mucosa in CD [6].
59 Genome wide association studies (GWAS) have identified 240 IBD susceptibility loci to date

1
2
3 60 [7], and have confirmed association with previously recognised susceptibility genes including
4
5 61 *Nucleotide-binding oligomerisation domain-containing protein 2 (NOD2)*. GWAS have also
6
7 62 identified the strong association of CD with genes involved in the autophagy pathway,
8
9 63 including *autophagy-related protein (ATG)16L1*, *Immunity-related GTPase family M protein*
10
11 64 (*IRGM*) and *leucine rich repeat kinase 2 (LRRK2)* [8]. The strong association of IBD with
12
13 65 endoplasmic reticulum (ER) stress/Unfolded protein response (UPR) genes including *x-box-*
14
15 66 *binding protein 1 (XBP1)* [9] and genes involved in intestinal barrier function such as *MUC2*
16
17 67 [10] and *Anterior gradient 2 (AGR2)* [11] have been detected by gene targeted approaches.
18
19
20 68 Together, these genetic studies have led to increased research exploring links between
21
22
23 69 autophagy and ER stress/UPR dysregulation and IBD pathogenesis.
24
25
26

27 70 Autophagy

28
29 71
30
31
32 72 Autophagy is an intracellular process that plays an important housekeeping role by degrading
33
34 73 excessive, damaged or aged proteins and organelles to maintain cellular homeostasis [12].
35
36
37 74 Basal autophagy is tightly regulated by the coordinated activity of autophagy-related (ATG)
38
39 75 proteins [13] and constitutes an important survival mechanism induced in response to
40
41
42 76 multiple stress conditions such as nutrient deprivation, hypoxia, DNA damage or intracellular
43
44 77 pathogens [12]. There are three main types of autophagy in mammalian cells;
45
46 78 macroautophagy (herein referred to as autophagy), microautophagy and chaperone-
47
48 79 mediated autophagy [12].
49
50
51
52
53

54 80 When autophagy is initiated a double membrane vesicle is formed (the autophagosome)
55
56 81 around the cargo to be degraded (**Figure 1**). The mature autophagosome then fuses with a
57
58
59 82 lysosome to form an autophagolysosome, in which lysosomal enzymes degrade the inner
60
61
62
63
64
65

1
2
3 83 membrane and cargo and the resulting macromolecules are released into the cytosol for
4
5
6 84 recycling (**Figure 1**).

7
8
9 85 Selective types of autophagy also exist, including autophagy of microorganisms (xenophagy)
10
11 86 and autophagy of the ER membrane (ER-phagy), which use specific receptors and adaptor
12
13 87 proteins to link the cargo to the autophagy machinery [14]. For example, Sequestosome
14
15 88 1/p62-like receptors (SLRs) target cytosolic pathogens and other cargo to initiate autophagy
16
17 89 [15]. SLRs function by binding to the small regulatory protein ubiquitin on the surface of cargo
18
19 90 [16–18] and subsequently associate with the autophagy machinery via a binding motif called
20
21 91 the LC3-interacting region (LIR) [19]. Adaptor proteins, such as autophagy-linked FYVE protein
22
23 92 (ALFY), can also bind ubiquitinated pathogens via p62 to promote association with the
24
25 93 autophagy machinery [20]. To date, five main types of SLR have been described;
26
27 94 sequestosome 1/p62, optineurin [18], NBR1 (Neighbor of BRCA1 gene 1) [21], NDP52 (Nuclear
28
29 95 Domain 10 Protein 52) [17] and the NDP52-like receptor calcoco3 (Calcium-binding and
30
31 96 coiled-coil domain-containing protein 3) [22], and specific cargo receptors are important for
32
33 97 distinct types of selective autophagy. For example, a recent study has shown that the non-
34
35 98 canonical cargo receptor cell-cycle progression gene 1 (CCPG1) is essential for ER-phagy [23],
36
37 99 while another study demonstrated an integral role for optineurin in the maintenance of ER
38
39
40
41
42
43
44
45 100 homeostasis by assisting the removal of hyper-activated UPR kinases [24].
46
47
48

49 101 Autophagy and CD

50
51
52
53

54 102 Autophagy affects many essential cellular processes and dysregulation of autophagy has been
55
56 103 linked to a multitude of human diseases [25]. Autophagy plays an important role in both
57
58
59 104 innate and adaptive immune signalling pathways and loss of immune regulation is a key event
60
61
62
63
64
65

105 leading to the chronic inflammation observed in CD [26]. Impaired autophagy responses have
106 been observed in a range of cell types derived from CD patients including the specialized
107 intestinal epithelial cells (IECs) Paneth cells and goblet cells, and leukocytes, such as
108 macrophages and dendritic cells [27].

109 Functional studies have linked impaired autophagy to CD-associated genetic variants in
110 *NOD2*, *ATG16L1*, *IRGM* and *LRRK2*. The single nucleotide polymorphism (SNP) in *ATG16L1*
111 causes a single amino acid change from threonine to alanine at position 300 (*T300A*) [28],
112 which is associated with Paneth cell and goblet cell dysfunction, and significantly impairs
113 autophagic clearance of pathogens [29–32]. *IRGM* is required for the initiation of xenophagy
114 and the clearance of intracellular organisms such as *Mycobacterium tuberculosis* [33] and
115 dysregulation of *IRGM* expression compromises the control of intracellular replication of CD-
116 associated adherent invasive *Escherichia coli* (AIEC) by autophagy [34]. *LRRK2* expression is
117 increased in colonic biopsy specimens from patients with CD [35] and functionally *LRRK2* can
118 enhance NFκB-dependent transcription, while small interfering RNA [siRNA] knockdown of
119 *LRRK2* interferes with bacterial killing [35].

120 *NOD2* is a member of the Nod-like receptor (NLR) family of pattern recognition receptors
121 (PRR) and recognises a component of the bacterial cell wall muramyl dipeptide (MDP) to
122 induce innate immune responses [36]. CD-associated *NOD2* SNPs (R702W, G908R and
123 L1007fs) affect the leucine rich repeat domain disrupting interaction with MDP and
124 abrogating immune responses initiated by this receptor [37]. The immunoregulatory
125 properties of *NOD2* have also been linked to autophagy, and CD susceptibility is heightened
126 when *ATG16L1* and *NOD2* variants present in combination, causing synergistic genetic
127 epistasis [38,39]. A direct functional interaction between these proteins has been

128 determined; NOD2 was shown to recruit ATG16L1 to the plasma membrane to initiate
129 autophagy at the sites of bacterial entry [40], and in a separate study IRGM was shown to
130 regulate the formation of a complex containing NOD2 and ATG16L1 that is necessary for the
131 induction of xenophagy [41]. The interaction of IRGM with NOD2 also stimulates
132 phosphorylation cascades involving AMPK, ULK1 and Beclin1 that regulate autophagy
133 initiation complexes [41]. Cells harbouring CD-associated *NOD2* variants and/or the *ATG16L1*
134 *T300A* variant exhibit a number of disrupted functions linked to autophagy including reduced
135 production of antimicrobial peptides, enhanced pro-inflammatory responses and aberrant
136 activation of adaptive immune responses [40,42–44].

137 Significantly, abnormalities in the secretory capacity of Paneth cells are observed in mice
138 deficient for ATG16L1 [30,45,46], NOD2 [47,48], IRGM [49] and LRRK2 [50] indicating that
139 autophagy plays an essential and specific role in Paneth cell function. Despite the significant
140 effects on Paneth cell function, mouse strains developed for deficiency in functional ATG16L1
141 do not exhibit spontaneous intestinal inflammation [29–31]. In contrast, a mouse strain with
142 targeted deletion of *ATG16L1* in IECs developed a spontaneous transmural ileitis similar to
143 ileal CD [24]. Furthermore, targeted deletion of *ATG16L* in haematopoietic cells can enhance
144 susceptibility to DSS-induced acute intestinal injury in mice [51] and ATG16L1 deficiency in
145 myeloid cells in a mouse strain led to disrupted macrophage function and bacterial clearance
146 [52]. Murine models with non-functional NOD2 do not develop spontaneous colitis [53],
147 however a *NOD2* mutation similar to the L1007fs mutation increased susceptibility to DSS-
148 induced colitis in mice [54]. *Irgm1*-deficient mice also exhibit abnormalities in Paneth cells,
149 accompanied by increased susceptibility to inflammation in the colon and ileum [49]. Finally,
150 *LRRK2* deficiency confers enhanced susceptibility to experimental colitis in mice, which was

151 associated with enhanced nuclear localisation of the transcription factor nuclear factor of
152 activated T cells (NFAT1), important for regulating innate immune responses [55].

153 ER-stress and UPR signalling

154 ER stress results from accumulation of unfolded and misfolded protein in the ER, and the UPR
155 is activated to resolve ER stress and restore homeostasis. The UPR inhibits protein synthesis,
156 promotes protein re-folding, and induces degradation of unfolded and misfolded proteins
157 through ER-associated protein degradation (ERAD) and autophagy (**Figure 2**). If these survival
158 mechanisms are unsuccessful, the UPR can induce apoptosis [56]. The major regulators of the
159 UPR are the ER-membrane resident proteins PERK (protein kinase RNA-like endoplasmic
160 reticulum kinase), inositol-requiring transmembrane kinase endonuclease 1 (IRE1) and
161 activated transcription factor (ATF)6. When inactive these proteins are bound to binding
162 immunoglobulin protein (BiP), also known as glucose regulated protein 78 (GRP78) [57].
163 During ER stress, BiP binds to misfolded proteins in the ER and dissociates from the ER-
164 membrane resident proteins to allow their transition to an active state [57] (**Figure 2**).

165 When active, PERK phosphorylates elongation initiation factor 2 α (EIF2 α), to inhibit general
166 protein synthesis [58] and specifically up-regulates ATF4 [59]. ATF4 in turn transcriptionally
167 up-regulates several other UPR genes including CCAAT/enhancer-binding protein (C/EBP)
168 homologous protein (CHOP) [60,61] (**Figure 2**). CHOP is also a transcription factor that
169 regulates several other UPR genes, and under conditions of prolonged ER stress can promote
170 apoptosis [60,61].

171 IRE1 exists in two forms: IRE1 α that is ubiquitously expressed and IRE1 β that is only expressed
172 in the GI tract and lung epithelial cells [62]. During ER stress, IRE1 is activated through

173 dimerization and auto-phosphorylation [63,64]. The IRE1 α RNase domain is essential for
174 creating transcriptionally activate *XBP1* messenger RNA (mRNA) via splicing, which acts as a
175 transactivator of UPR genes [65–68] (**Figure 2**). IRE1 endoribonuclease activity also facilitates
176 degradation of specific mRNA in a process known as RIDD (regulated IRE1-dependent decay)
177 [69].

178 ATF6 translocates to the Golgi apparatus once released from its complex with BiP [70]. This
179 allows cleavage by site 1 and site 2 proteases (S1P and S2P), which releases the
180 transcriptionally active cytoplasmic domain of ATF6 (ATF6-N) that induces UPR-associated
181 genes [71–73] (**Figure 2**). Among the ATF6 upregulated genes are *CHOP* and *XBP1* [74].

182 ER-stress, UPR and intestinal inflammation

183 Genetic studies have identified several ER-stress/UPR genes associated with IBD [75].
184 Moreover, ER-stress levels are increased in ileal and colonic biopsies from CD patients, with
185 higher than normal levels of BiP, chaperone protein Gp96, and spliced *XBP1* observed [9,76–
186 78] (**Table 1**). Several studies have focused on IRE1-XBP1 signalling in murine models. In mice
187 with targeted deletion of XBP1 in intestinal epithelial cells (IECs) (*XBP1 Δ IEC* mice), spontaneous
188 inflammation of the small intestine, increased susceptibility to DSS-induced colitis and
189 elevated levels of ER stress were observed [9] (**Table 1**). Furthermore, in *XBP1 Δ IEC* mice
190 increased levels of apoptosis were observed along with reduced goblet cell and Paneth cell
191 numbers, leading to decreased production of host defence peptides and higher susceptibility
192 to *Listeria monocytogenes* infection [9] (**Table 1**). *XBP1* has also been shown to suppress
193 experimental colitis-associated cancer [79], and is essential for efficient TLR-mediated pro-

194 inflammatory responses to infection in macrophages [80]. These studies support *XBP1* as a
195 key component of the protective function of IECs and macrophages.

196 Although the UPR acts to maintain ER-homeostasis, hyper-activation of certain UPR
197 components can create a pro-inflammatory state. In *XBP1^{ΔIEC}* mice increased activation of
198 IRE1 α , causes hyper activation of NF κ B, and spontaneous inflammation [45] (**Table 1**). IRE1 β
199 knock-out mice have enhanced sensitivity to DSS-induced colitis [81] and exhibit goblet cell
200 abnormalities with exaggerated MUC2 accumulation (**Table 1**). In contrast, IRE1 α knock-out
201 mice have normal goblet cells [82]. In murine Paneth cells, IRE1 α and IRE1 β have distinct roles
202 with hyper activation of IRE1 α driving CD-like ileitis, and IRE1 β having a protective role [24].

203 Association of aberrant PERK-EIF2 α and ATF6 pathways with intestinal inflammation have
204 also been identified. A mouse model expressing non-phosphorylatable EIF2 α in IECs resulted
205 in functional abnormalities in Paneth cells and increased susceptibility to *Salmonella* infection
206 and DSS-induced colitis [83] (**Table 1**). ATF6 α deficient mice exhibit increased ER stress as
207 indicated by elevated levels of BiP, ATF4, CHOP and spliced *XBP1*, which result in enhanced
208 sensitivity to DSS-induced colitis [84] (**Table 1**). Additionally, hypomorphic mutation in
209 *membrane-bound transcription factor peptidase S1P-encoding gene (Mbtps1)*, which encodes
210 the S1P responsible for cleavage of ATF6, causes enhanced susceptibility to DSS-induced
211 colitis [85]. Although there is less evidence to support a role for PERK-EIF2 α and ATF6
212 pathways in IBD pathogenesis, their importance for ER stress responses in the intestinal
213 epithelium is clear.

214 ER-stress and intestinal barrier function

215 In the intestinal epithelium, cells that naturally secrete large amounts of protein, such as
1
2
3 216 Paneth cells and goblet cells, are more susceptible to ER-stress and therefore rely heavily on
4
5 217 the UPR to maintain homeostasis. MUC2 is the major component of mucin that is produced
6
7
8 218 in goblet cells and secreted into the intestinal lumen. *Winnie* mice are characterised by a
9
10 219 missense mutation in *MUC2*, which causes abnormalities in goblet cells, leading to aberrant
11
12
13 220 mucous production and spontaneous colitis, and association with *MUC2* variants has been
14
15 221 identified in IBD patients [86]. *Winnie* mice also exhibit severe ER stress in goblet cells [10],
16
17
18 222 which causes up to four-fold increase in activated dendritic cells in the colonic lamina propria,
19
20
21 223 and aberrant adaptive immune responses associated with interleukin (IL)-23/Th17 [87].
22
23 224 Goblet cell abnormalities are also apparent in mice deficient in UPR transcription factor
24
25
26 225 OASIS, which causes increased ER stress and susceptibility to DSS-induced colitis [88,89].
27
28
29 226 AGR2 is an ER resident protein highly expressed in goblet and Paneth cells and regulates the
30
31 227 formation of disulphide bonds in mature proteins. *AGR2*^{-/-} mice exhibit a decreased number
32
33
34 228 of goblet cells and MUC2 production, Paneth cell abnormalities, elevated ER-stress and
35
36 229 spontaneous colitis [90]. Notably, AGR2 is decreased in patients with CD and UC [11]. These
37
38
39 230 studies highlight the key role of intestinal secretory cells and breakdown of intestinal barrier
40
41 231 function in IBD pathogenesis.
42
43
44

232 Functional intersection between autophagy and the UPR

45
46
47
48
49

50 233 The UPR and autophagy are intimately linked processes. In a range of Intestinal epithelial cells,
51
52
53 234 chemical ER-stress inducers activate autophagy, modulated by enhanced expression of CHOP
54
55 235 and stimulation of the IRE1 α pathway [91]. In endothelial cells, IRE1 α -dependent splicing of
56
57
58 236 *XBP1* mRNA activated autophagy via up-regulation of *Beclin-1*, which is a major regulator of
59
60
61
62
63
64
65

237 the autophagy pathway [92] (**Figure 3**). Contrary to expectation, *XBP1* deletion in a familial
238 amyotrophic lateral sclerosis mouse model increased autophagy, which enhanced clearance
239 of accumulated toxic superoxide dismutase-1 (SOD1) aggregates [93]. It was suggested that
240 in this scenario, autophagy is induced in a compensatory manner due to attenuated UPR.

241 The UPR and autophagy also intersect at the PERK-EIF2 α -ATF4 pathway [94–99]. In an *in vitro*
242 model of osteosarcoma, PERK induced autophagy via mechanistic target of rapamycin
243 (mTORC1) inhibition to promote survival in response to ER stress-conferred chemoresistance
244 to apoptosis [95] (**Figure 3**). Additionally, PERK modulates autophagy via AMPK-dependent
245 inhibition of mTORC1 in response to extracellular matrix (ECM) detachment in mammary
246 epithelial cells (MECs) [94]. One of the main functional outcomes of PERK signalling is reduced
247 protein synthesis. Inhibition of mTORC1 helps to promote this effect as mTORC1 controls
248 synthesis of ~15-20% of protein within the cell [100]. Thus, via modulation of mTORC1, PERK
249 signalling achieves dual outcomes; inhibition of protein synthesis and induction of autophagy
250 to degrade misfolded proteins.

251 During amino acid deprivation, ATF4 and CHOP can bind specific C/EBP-ATF Response
252 Elements (CAREs), also known as Amino Acid Response Elements (AAREs) and CHOP-Response
253 Elements (CHOP-REs) to induce transcription of a wide range of autophagy genes [101] (**Figure**
254 **3**). In other studies, hypoxia or ECM detachment induced PERK-dependent autophagy due to
255 autophagy gene up-regulation via ATF4 and CHOP [102–104]. This up-regulation of autophagy
256 gene transcription by the UPR was shown to replenish autophagy proteins to promote survival
257 during cellular stress [103].

258 AT6 has also been implicated mechanistically in autophagy regulation. In response to cellular
1
2
3 259 stress, interferon (IFN)- γ activates the Ask1 (Apoptosis signal-regulating kinase 1)/MAPK
4
5 260 (Mitogen-activated protein kinase) pathway, which phosphorylates AT6 to allow its
6
7
8 261 proteolytic activation [105]. AT6 interaction with C/EBP- β is essential for IFN- γ -induced up-
9
10 262 regulation of *DAPK1* (*death-associated protein kinase 1*), which can subsequently stimulate
11
12
13 263 autophagy [106] (**Figure 3**). Mice lacking either AT6 or Ask1 are highly susceptible to bacterial
14
15 264 infection due to defective autophagy [105,106]. Furthermore, AT6 recruitment of DAPK1 in
16
17
18 265 response to ER stress enhanced xenophagy in human colonic biopsies and epithelial cells,
19
20
21 266 which was attenuated in cells harbouring the *ATG16L1 T300A* SNP [107]. Additionally,
22
23 267 activated AT6 was shown to stimulate Akt (protein kinase B), which resulted in the inhibition
24
25
26 268 of mTORC1 [108,109] (**Figure 3**).

27
28
29 269 In a recent study in MCF-7 human breast cancer cells, ER stress induced by the chemo-
30
31
32 270 preventative agent ursolic acid (UA) was associated with autophagy activation [99]. UA
33
34
35 271 induced autophagy via MAPK1/3 signalling and subsequent promotion of PERK signalling,
36
37
38 272 resulting in the inhibition of apoptosis. Furthermore, a study in human ovarian cancer cells
39
40 273 showed interdependent activation of autophagy and the PERK-EIF2 α UPR pathway when
41
42
43 274 treated with metformin, which causes energy starvation [98]. In these scenarios an
44
45 275 unconventional relationship between autophagy and ER stress was uncovered, which remains
46
47
48 276 to be mechanistically solved. Nonetheless, under these circumstances the interaction of the
49
50 277 UPR and autophagy pathways has pro-survival outcomes.

51
52
53
54 278 **Convergence of autophagy, ER-stress and CD**
55
56
57
58
59
60
61
62
63
64
65

279 In an attempt to relieve ER-stress the UPR can induce autophagy to degrade misfolded
1
2
3 280 proteins, protein aggregates and damaged organelles [91,110–113]. Autophagy activity is
4
5 281 increased in highly secretory Paneth cells [45] to counterbalance high levels of ER-stress
6
7 282 [112], thus ER-stress is a significant risk in these cells when the UPR or autophagy is not
8
9
10 283 functional. Consistent with this, in Paneth cells of CD patients harbouring *ATG16L1 T300A* risk
11
12 284 alleles, BiP and pEIF2 α are highly expressed [46] (**Table 1**). Significantly, *ATG16L1;XBP1 ^{Δ IEC}*
13
14 285 mice develop similar phenotypic ileitis to *ATG16L1 ^{Δ IEC}* mice, but earlier in life due to increased
15
16
17 286 ER stress [24,45].

22 287 ERAD can regulate the degradation of IRE1 α to prevent accumulation of toxic IRE1 α
23
24 288 aggregates, however persistent ER-stress will inhibit ERAD degradation of IRE1 α [24]. When
25
26
27 289 this occurs, autophagy plays an important role in the clearance of supramolecular clusters of
28
29
30 290 IRE1 α (**Figure 3**). In *ATG16L1 ^{Δ IEC}* mice, development of spontaneous CD-like ileitis is associated
31
32 291 with defective autophagy resulting in toxic accumulation of IRE1 α in Paneth cells [24] (**Table**
33
34
35 292 **1**). Furthermore, the selective autophagy receptor optineurin interacts with IRE1 α , and
36
37
38 293 optineurin deficiency amplified the accumulation of IRE1 α [24]. In humans homozygous for
39
40 294 *ATG16L1 T300A*, a similar accumulation of IRE1 α was observed in intestinal epithelial crypts
41
42
43 295 [24] (**Table 1**). This has led to suggestion that the *ATG16L1 T300A* SNP may define a specific
44
45
46 296 subtype of patients with CD, characterised by Paneth cell ER-stress [46]. This synergistic and
47
48 297 compensatory relationship between the UPR and autophagy is affirmed by the presence of
49
50
51 298 CD-associated SNPs in *ATG16L1* and *XBP1*.

54 299 A recent study has demonstrated a direct link between NOD1/2 and the IRE1 α pathway in the
55
56
57 300 context of ER-stress-induced inflammation [114]. When active, IRE1 α stimulates the c-Jun N-
58
59
60 301 terminal kinase (JNK) pathway and recruits TRAF2 (TNF receptor-associated factor 2) to the
61
62
63
64
65

302 ER membrane to trigger NF κ B signalling [115,116] and autophagy induction [112,117,118]
1
2
3 303 **(Figure 3)**. In mouse and human cells, ER-stress induced by chemicals or infection with
4
5 304 *Brucella abortus* and *Chlamydia muridarum* increased inflammation and IL-6 production
6
7 305 [114]. This response was dependent on NOD1/2 and receptor-interacting serine/threonine-
8
9 306 protein kinase 2 (RIPK2), but also on IRE1 α kinase activity and TRAF2-induced NF κ B signalling
10
11 307 [114]. This suggests there is a functional intersection between the IRE1 α pathway and
12
13 308 NOD1/2 signalling, which is facilitated by TRAF2 **(Figure 3)**.
14
15
16
17
18

19 309 Interestingly, an additional study has shown that ER-stress responses can be modulated by
20
21 310 another innate immune sensor called stimulator of interferon genes (STING) in response to
22
23 311 cyclic-di-AMP (c-di-AMP), a vita-PAMP (pathogen associated molecular pattern) present in
24
25 312 live Gram-positive bacteria [119]. This process induces autophagy via inhibition of the major
26
27 313 autophagy suppressor mTORC1 and localisation of STING to autophagosomes.
28
29
30
31
32

34 314 Pharmacological induction of autophagy and the UPR

35
36
37

38 315 A recent review estimated IBD treatment costs of £720 million (\$940m) per year in the United
39
40 316 Kingdom alone [120], with roughly a quarter of these costs directly attributed to drug
41
42 317 treatments [121]. The efficacy of these drugs continues to come under scrutiny as response
43
44 318 to treatment often diminishes over time, with a review of worldwide cohorts estimating that
45
46 319 between 10–35% of CD patients required surgery within a year of diagnosis and up to 61% by
47
48 320 10 years [122]. In order to improve the efficacy of IBD treatment, optimization of existing
49
50 321 clinical therapies and the development of novel therapeutics is required.
51
52
53
54
55
56

57 322 The convergence between autophagy and UPR pathways provides new opportunity for the
58
59 323 treatment of IBD and the modulation of the UPR in combination with autophagy inducers is a
60
61
62
63
64
65

1 324 promising therapeutic strategy. There is evidence that inducing autophagy can have
2
3 325 therapeutic benefits for the treatment of IBD [26] with several studies investigating the utility
4
5 326 of autophagy inducers as adjuvant therapies. Rapamycin analogues, sirolimus and everolimus,
6
7 327 inhibit mTORC1 to induce autophagy and are already approved for clinical use for post-
8
9
10 328 transplantation (e.g. liver and renal) management. In IL-10-deficient mice, everolimus
11
12
13 329 treatment alleviated spontaneous colitis and reduced CD4+ T cells and IFN- γ [123]. In a case
14
15 330 study sirolimus improved symptoms and intestinal healing in a patient with severe refractory
16
17
18 331 CD [124]. In another case study, symptoms were controlled for 18 months with everolimus
19
20
21 332 treatment in a refractory UC patient [125]. Moreover, in a study of refractory paediatric IBD,
22
23 333 sirolimus induced clinical remission in 45% of UC patients and 100% of CD patients; albeit the
24
25
26 334 sample size was small [126]. Significantly, everolimus had comparable safety and tolerability
27
28
29 335 as azathioprine when used to maintain steroid-induced remission in a cohort of adult CD
30
31 336 patients [127]. As these mTORC1 inhibitors are already approved for clinical use, they have
32
33
34 337 been investigated the most extensively, however there are a plethora of novel autophagy
35
36 338 modulators that are currently being developed, characterised and patented for therapeutic
37
38
39 339 use in a range of diseases including IBD [128,129].

40
41
42 340 Recent progress has also been made to identify specific chemical inducers of the UPR. A
43
44
45 341 screen of 1,200 FDA-approved compounds carried out in *C.elegans* identified eight
46
47
48 342 compounds that induced UPR responses, four of which specifically increased mitochondrial
49
50 343 UPR [130]. The identified drugs included antirheumatic agents, antianginal calcium channel
51
52
53 344 blockers; androgen receptor inhibitors used for cancer therapy and tetracycline antibiotics.

54
55
56 345 A well-characterised modulator of the UPR, tauroursodeoxycholic acid (TUDCA), that
57
58
59 346 promotes protein refolding to reduce ER-stress, was shown to ameliorate DSS-induced colitis
60
61
62
63
64
65

1 347 in mice by decreasing ER-stress in IECs [84]. Furthermore, a selective inhibitor of eIF2 α
2
3 348 dephosphorylation protects cells from ER-stress and ameliorates murine experimental colitis
4
5 349 [131,132]. Supplementation with glutamine has also been suggested for the improvement of
6
7 350 IBD treatment, as this amino acid was shown to dampen experimental colitis in rats by
8
9
10 351 inhibiting ER-stress in colonic epithelial cells [133].
11

12
13
14 352 Drugs used to treat metabolic disorders have also been investigated for UPR inducing
15
16 353 properties. The biguanides metformin and phenformin have been implicated in induction of
17
18
19 354 the UPR and resolution of ER-stress via activation of AMPK, which subsequently stimulated
20
21
22 355 IRE1 α and PERK pathways [98,134,135]. Inhibitors of dipeptidyl peptidase IV (DPP4), including
23
24 356 gemigliptin, also prevented ER-stress-mediated apoptosis by promoting IRE1 α and PERK
25
26
27 357 pathways [136]. Furthermore, agonists of the glucagon-like peptide-1 receptor, such as
28
29
30 358 exenatide, relieved ER stress via up-regulation of *ATF4* expression [137]. Exogenous chemical
31
32 359 chaperones have also been explored as a method to relieve ER stress by mimicking ER
33
34
35 360 chaperones to promote protein transport and re-folding capacity [138].
36

37
38 361 Although several studies have demonstrated beneficial effects of enhancing UPR function for
39
40
41 362 intestinal homeostasis, future investigations should proceed with caution. For example,
42
43
44 363 hyper-activation of the UPR kinase IRE1 α can exacerbate intestinal inflammation, as seen in
45
46 364 patients with *ATG16L1* and *NOD2* mutations, therefore, in certain circumstances
47
48
49 365 pharmacological inhibition of UPR receptors would be a more effective strategy [24,45,114]
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

366 Of particular interest, the selective autophagy cargo receptor optineurin forms a critical link
1
2
3 367 between ER-stress resolution and autophagy due to its role in the degradation of IRE1 α
4
5 368 aggregates [24], and another recently identified autophagy cargo receptor that is integral for
6
7
8 369 resolution of ER-stress, CCPG1, mediates ER-phagy to remove damaged ER membranes [23].
9
10 370 Understanding the biology and functions of adaptors such as optineurin and CCPG1 may
11
12
13 371 identify novel druggable targets and expedite development of the next generation of
14
15
16 372 therapeutics aimed at modulation of the UPR in combination with autophagy.
17
18

19 373 Discussion

20
21
22

23 374 The complexity of IBD is evident from the large number of risk loci identified by genetic
24
25
26 375 studies, and the diverse health profile of patients that are affected. Mouse models of IBD
27
28
29 376 cannot emulate the human disease, however they are useful tools to explore how specific
30
31
32 377 gene mutations influence inflammation. Interestingly, as highlighted in **(Table 1)** the majority
33
34 378 of mouse models mimicking IBD-associated genetic risk do not develop spontaneous
35
36
37 379 inflammation, but rather they are sensitised to DSS-induced colitis, which acts by damaging
38
39 380 the epithelium and increasing intestinal permeability. The intestinal epithelium has
40
41
42 381 important immunoregulatory functions and controls the equilibrium between tolerance and
43
44 382 immunity to non-self-antigens [139]. As such breakdown of intestinal epithelial barrier
45
46
47 383 function and concomitant interaction with environmental factors in the lumen is a trigger for
48
49
50 384 inflammation. The intestinal lumen comprises a multitude of potential triggers including the
51
52 385 microbiota, dietary antigens, and luminal antigens. Additional triggers may be host-derived
53
54
55 386 factors that are released into the lumen as the intestinal epithelial barrier breaks down. These
56
57 387 so-called Damage-Associated Molecular Patterns (DAMPS) include intracellular proteins, such
58
59
60 388 as high-mobility group box 1 (HMGB1), heat-shock proteins and components derived from
61
62
63
64
65

1 389 the extracellular matrix. Examples of non-protein DAMPs include genomic DNA,
2
3 390 mitochondrial DNA, RNA, uric acid and ATP [140,141]. Not surprisingly, there is considerable
4
5 391 interest in developing novel therapeutic strategies aimed at re-establishing intestinal barrier
6
7 392 function [142] and modulation of DAMPs for the treatment of IBD [140].
8
9

10
11 393 Dysbiosis of the gut microbiome is strongly implicated in the pathogenesis of CD [143], and it
12
13
14 394 has been suggested that microbial dysbiosis may be an environmental trigger. A recent study
15
16 395 by Tschurtschenthaler and colleagues [24] addressed this question. Although microbial
17
18
19 396 dysbiosis was present in the ileum of *Atg16l1;Xbp1*^{ΔIEC} mice, such structural alteration of the
20
21
22 397 microbiota did not trigger ileitis but, rather, aggravated DSS-induced colitis [24]. In order to
23
24 398 understand the role of the environment in disease, determining the relative contribution of
25
26
27 399 genetics and a detailed characterization of environmental triggers is required.
28
29

30
31 400 Greater understanding of the genetic factors that underlie CD pathogenesis are leading to
32
33 401 improvements in treatment. Development of personalised therapies may be achieved via
34
35
36 402 genotyping for key SNPs in genes involved in both the autophagy and UPR pathways. IBD
37
38
39 403 drugs already established in the clinic have been shown to exert their effects, at least in-part,
40
41 404 through the modulation of autophagy [26] or the UPR, and establishing patient genotypes
42
43
44 405 may help predict response. For example, recent studies have identified an association
45
46 406 between *ATG16L1 T300A* SNP and an enhanced therapeutic effect of thiopurines [144] and
47
48
49 407 anti-TNF- α therapy [145]. Interestingly, the immunoregulatory effects of these drugs were
50
51 408 associated with autophagy stimulation [144,146,147] and the *T300A* genotype has been
52
53
54 409 associated with a subset of patients that exhibit deficiencies in both the UPR and autophagy
55
56
57 410 [46]. Furthermore, CD patients harbouring *NOD2* mutations associate with better clinical
58
59 411 outcomes in response to thiopurines, whereas CD patients with wild-type *NOD2* respond
60
61
62
63
64
65

1
2
3 412 better to steroids and anti-TNF therapy [148]. Due to the genetic complexity of IBD and
4
5 413 epistasis between genes, it is imperative that multiple genes are analysed for the purpose of
6
7 414 patient stratification. For example, a recent study identified a 32-gene transcriptomic
8
9 415 signature in lymphoblastoid cells that was able to predict lack of response to thiopurines, with
10
11 416 aberrant cell cycle control, DNA mismatched repair and RAC1-dependent mechanisms
12
13 417 implicated in thiopurine resistance [149]. Furthermore, it is increasingly clear that epigenetic,
14
15 418 microRNA and immune cell signatures among others will have a significant role to play in
16
17 419 predicting disease susceptibility and response to therapy [150–152].
18
19
20
21

22 420 With regards to the intestinal microbiota, a recent study has characterised microbial
23
24 421 signatures for the diagnosis of IBD that were highly sensitive and could differentiate CD
25
26 422 patients from healthy controls and UC patients. This study highlights the potential for using
27
28 423 the intestinal microbiota as a micro-biomarker [153]. Importantly, as many drugs need to be
29
30 424 metabolised and de-toxified by the gut microbiota, this approach could also have application
31
32 425 in predicting response to therapy. Given that dysregulation of autophagy and ER-stress can
33
34 426 affect the intestinal microbial environment, analysis of microbial signatures may help to
35
36 427 determine if a patient would benefit from drugs that modulate the autophagy or UPR
37
38 428 pathways.
39
40
41
42
43
44
45

46 429 To conclude, the ER-stress/UPR and autophagy pathways play a vital role in the maintenance
47
48 430 of intestinal homeostasis and breakdown of these converging pathways has been implicated
49
50 431 in persistent intestinal infections, chronic inflammation and dysregulated immune responses
51
52 432 observed in IBD. Therefore, strategies aimed at modulating these pathways simultaneously
53
54 433 may prove to be an effective therapeutic option.
55
56
57
58
59
60
61
62
63
64
65

1 **434** Funding

2
3
4 **435** This work was supported by a Crohn's in Childhood Research Association (CICRA) PhD
5
6
7 **436** studentship to KMH. PH is supported by a NHS Research Scotland Career Researcher
8
9
10 **437** Fellowship.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

438 Figure Legends

439 Figure 1: Autophagy pathway and autophagosome biogenesis

440 During the initial stages of autophagy, the isolation membrane forms a double membrane
441 vesicle (the autophagosome) around the cargo to be degraded. ULK complex (ULK1-ULK2-
442 ATG13-FIP200-ATG101) and Beclin 1 (Vps34-Vps150-Beclin1) complex, through interaction
443 with ATG14, recruit autophagy proteins and complexes to the autophagosome membrane.
444 ATG12 is conjugated to ATG5 and forms a complex with ATG16L1 (ATG16L1 complex). The
445 ATG16L1 complex is proposed to specify the site of LC3 lipidation for autophagosome
446 formation. LC3 is conjugated to PE to form lipidated LC3-II and is associated with the
447 autophagosome outer membrane. Upon autophagosome closure, LC3 localises to the inner
448 membrane and other autophagy proteins and complexes dissociate for recycling. The mature
449 autophagosome then fuses with a lysosome to form an autophagolysosome, in which cargo
450 are degraded by lysosomal enzymes and subunits are recycled.

451 Figure 2: The unfolded protein response

452 BiP chaperone protein binds unfolded/misfolded proteins in the ER and dissociates from
453 transmembrane receptors upon accumulation of the toxic proteins. The transmembrane
454 receptors PERK, IRE1 α and ATF6 become activated. PERK phosphorylates EIF2 α , which
455 downregulates global translation but specifically upregulates ATF4 and CHOP that upregulate
456 UPR-associated genes. IRE1 α splices XBP1 to its active form and ATF6 is cleaved by S1P and
457 S2P to active ATF6-N, which both translocate to the nucleus to upregulate UPR-associated
458 genes. The main function of these UPR-associated genes is to increase protein refolding,

1
2
3 459 inhibit synthesis of new protein and degrade unfolded/misfolded proteins through autophagy
4
5
6 460 and ERAD.

7
8
9
10 461 **Figure 3: Intersection between autophagy and the unfolded protein response**


11
12 462 ER stress activates transmembrane receptors PERK, IRE1 α and ATF6. PERK phosphorylates
13 463 EIF2 α , which specifically upregulates ATF4 and CHOP that bind AAREs and CHOP-Res to
14
15 464 upregulate autophagy genes. PERK also induces autophagy via mTORC1 inhibition. IRE1 α
16
17 465 splices XBP1 to its active form, which up-regulates *Beclin-1*. IRE1 α endonuclease activity
18
19 466 activates the JNK pathway, which induces autophagy via TRAF2, NOD2 and NF κ B. Enhanced
20
21 467 autophagy degrades accumulated IRE1 α clusters. Active ATF6-N induces autophagy via
22
23 468 mTORC1 inhibition and binds C/EBP- β to up-regulate *DAPK1*.

24
25
26
27
28 469 **Table 1: Murine models of intestinal inflammation**

29
30
31
32 470 Links between autophagy, ER-stress/UPR and experimental colitis/intestinal inflammation
33
34
35 471 and IBD.

36
37
38
39 472

40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65



473

474 Figure 1




Figure 2

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

475
476




Figure 3

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

477
478

Autophagy/ UPR pathway	Murine models of intestinal inflammation	IBD patients
ATG16L1	<ul style="list-style-type: none"> • ATG16L1 deficiency caused enhanced susceptibility to experimental colitis, Paneth cell and Goblet cell dysfunction, disrupted macrophage function and significantly impairs xenophagy [29-32, 51, 52] • ATG16L1 deletion in IECs induced spontaneous transmural ileitis [24] 	<i>ATG16L1 T300A</i> CD-associated SNP [28]
NOD2	NOD2 mutation causes enhanced susceptibility to DSS-induced colitis [54] and causes Paneth cell dysfunction [47, 48]	<i>NOD2</i> CD-associated SNPs (R702W, G908R and L1007fs) [37]
IRGM	<i>Irgm1</i> deficiency causes abnormalities in Paneth cells and increased susceptibility to inflammation in the colon and ileum [49]	<i>IRGM</i> CD-associated SNP [8]
LRRK4	LRRK2 deficiency confers enhanced susceptibility to experimental colitis in mice [55] and Paneth cell abnormalities [50]	<i>LRRK4</i> CD-associated SNP [8]
IRE1 α -XBP1	<ul style="list-style-type: none"> • <i>XBP1</i> deletion causes spontaneous intestinal inflammation, abnormal Paneth and goblet cell function and increased infection [9] • <i>XBP1</i> deletion causes overactivation of IRE1α and NFκB [45] • <i>ATG16L1</i> deletion causes accumulation of IRE1α in Paneth cells resulting in CD-like ileitis [24] 	<ul style="list-style-type: none"> • <i>XBP1</i> CD-associated SNP [9] • Increased levels of spliced <i>XBP1</i>, BiP and Gp96 in CD [9, 76-78] • <i>T300A</i> SNP causes accumulation of IRE1α in intestinal crypts [24]
IRE1 β	IRE1 β deletion causes enhanced sensitivity to DSS-colitis [81], goblet cell abnormalities and MUC2 accumulation [24]	
PERK-EIF2 α	Non-phosphorylatable EIF2 α caused Paneth cell abnormalities, enhanced DSS-colitis susceptibility and increased <i>Salmonella</i> infection [83]	Increased p-EIF2 α and BiP in CD patients with <i>T300A</i> SNP [46]
ATF6	<ul style="list-style-type: none"> • ATF6 deletion enhanced DSS-colitis susceptibility [84] • Mutation in <i>Mbtps1</i> (encodes S1P) causes enhanced DSS-colitis susceptibility [85] 	
AGR2	AGR2 deletion causes decreased Goblet cells and MUC2 production, Paneth cell abnormalities, elevated ER-stress and spontaneous colitis [90]	<ul style="list-style-type: none"> • <i>AGR2</i> CD-associated SNP [11] • <i>AGR2</i> decreased in IBD [11]

480 Table 1

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

481 References

- 1
2
3
4 482 1 Gasparetto M, Guariso G. Highlights in IBD Epidemiology and Its Natural History in the
5 Paediatric Age. *Gastroenterol Res Pr* 2013;**2013**:829040. doi:10.1155/2013/829040
6 483
7
8 484 2 Fakhoury M, Negrulj R, Mooranian A, *et al.* Inflammatory bowel disease: clinical aspects
9 485 and treatments. *J Inflamm Res* 2014;**7**:113–20. doi:10.2147/JIR.S65979
10
11 486 3 Neurath MF. Current and emerging therapeutic targets for IBD. *Nat Rev Gastroenterol*
12 *Hepatol* 2017;**14**:nrgastro.2016.208. doi:10.1038/nrgastro.2016.208
13 487
14
15 488 4 Boyapati R, Satsangi J, Ho GT. Pathogenesis of Crohn's disease. *F1000Prime Rep*
16 489 2015;**7**:44. doi:10.12703/p7-44
17
18
19 490 5 Zuo T, Ng SC. The Gut Microbiota in the Pathogenesis and Therapeutics of Inflammatory
20 491 Bowel Disease. *Front Microbiol* 2018;**9**:2247. doi:10.3389/fmicb.2018.02247
21
22 492 6 Darfeuille-Michaud A, Boudeau J, Bulois P, *et al.* High prevalence of adherent-invasive
23 493 *Escherichia coli* associated with ileal mucosa in Crohn's disease. *Gastroenterology*
24 494 2004;**127**:412–21. doi:10.1053/j.gastro.2004.04.061
25
26
27 495 7 Lange KM de, Moutsianas L, Lee JC, *et al.* Genome-wide association study implicates
28 immune activation of multiple integrin genes in inflammatory bowel disease. *Nat Genet*
29 496 2017;**49**:256–61. doi:10.1038/ng.3760
30 497
31
32 498 8 Franke A, McGovern DP, Barrett JC, *et al.* Genome-wide meta-analysis increases to 71 the
33 499 number of confirmed Crohn's disease susceptibility loci. *Nat Genet* 2010;**42**:1118–25.
34 500 doi:10.1038/ng.717
35
36
37 501 9 Kaser A, Lee A-H, Franke A, *et al.* XBP1 Links ER Stress to Intestinal Inflammation and
38 502 Confers Genetic Risk for Human Inflammatory Bowel Disease. *Cell* 2008;**134**:743–56.
39 503 doi:10.1016/j.cell.2008.07.021
40
41
42 504 10 Heazlewood CK, Cook MC, Eri R, *et al.* Aberrant Mucin Assembly in Mice Causes
43 505 Endoplasmic Reticulum Stress and Spontaneous Inflammation Resembling Ulcerative
44 506 Colitis. *PLoS Med* 2008;**5**. doi:10.1371/journal.pmed.0050054
45
46
47 507 11 Zheng W, Rosenstiel P, Huse K, *et al.* Evaluation of AGR2 and AGR3 as candidate genes for
48 508 inflammatory bowel disease. *Genes Immun* 2006;**7**:11–8. doi:10.1038/sj.gene.6364263
49
50
51 509 12 Dikic I, Elazar Z. Mechanism and medical implications of mammalian autophagy. *Nat Rev*
52 510 *Mol Cell Biol* 2018;**19**:349. doi:10.1038/s41580-018-0003-4
53
54 511 13 Lamb CA, Yoshimori T, Tooze SA. The autophagosome: origins unknown, biogenesis
55 512 complex. *Nat Rev Mol Cell Biol* 2013;**14**:759–74. doi:10.1038/nrm3696
56
57
58 513 14 Zaffagnini G, Martens S. Mechanisms of Selective Autophagy. *J Mol Biol* 2016;**428**:1714–
59 514 24. doi:10.1016/j.jmb.2016.02.004
60
61
62
63
64
65

- 515 15 Nys K, Agostinis P, Vermeire S. Autophagy: a new target or an old strategy for the
1 516 treatment of Crohn's disease? *Nat Rev Gastroenterol Hepatol* 2013;**10**:395–401.
2 517 doi:10.1038/nrgastro.2013.66
3
4
5 518 16 Dupont N, Lacas-Gervais S, Bertout J, *et al.* Shigella phagocytic vacuolar membrane
6 519 remnants participate in the cellular response to pathogen invasion and are regulated by
7 520 autophagy. *Cell Host Microbe* 2009;**6**:137–49. doi:10.1016/j.chom.2009.07.005
8
9
10 521 17 Thurston TLM, Ryzhakov G, Bloor S, *et al.* The TBK1 adaptor and autophagy receptor
11 522 NDP52 restricts the proliferation of ubiquitin-coated bacteria. *Nat Immunol*
12 523 2009;**10**:1215–21. doi:10.1038/ni.1800
13
14
15 524 18 Wild P, Farhan H, McEwan DG, *et al.* Phosphorylation of the autophagy receptor
16 525 optineurin restricts Salmonella growth. *Science* 2011;**333**:228–33.
17 526 doi:10.1126/science.1205405
18
19
20 527 19 Johansen T, Lamark T. Selective autophagy mediated by autophagic adapter proteins.
21 528 *Autophagy* 2011;**7**:279–96.
22
23 529 20 Filimonenko M, Isakson P, Finley KD, *et al.* The selective macroautophagic degradation of
24 530 aggregated proteins requires the PI3P-binding protein Alfy. *Mol Cell* 2010;**38**:265–79.
25 531 doi:10.1016/j.molcel.2010.04.007
26 532
27
28 532 21 Kirkin V, Lamark T, Sou Y-S, *et al.* A role for NBR1 in autophagosomal degradation of
29 533 ubiquitinated substrates. *Mol Cell* 2009;**33**:505–16. doi:10.1016/j.molcel.2009.01.020
30
31
32 534 22 Newman AC, Scholefield CL, Kemp AJ, *et al.* TBK1 kinase addiction in lung cancer cells is
33 535 mediated via autophagy of Tax1bp1/Ndp52 and non-canonical NF- κ B signalling. *PLoS One*
34 536 2012;**7**:e50672. doi:10.1371/journal.pone.0050672
35
36
37 537 23 Smith MD, Harley ME, Kemp AJ, *et al.* CCPG1 Is a Non-canonical Autophagy Cargo
38 538 Receptor Essential for ER-Phagy and Pancreatic ER Proteostasis. *Dev Cell* 2018;**44**:217-
39 539 232.e11. doi:10.1016/j.devcel.2017.11.024
40
41
42 540 24 Tschurtschenthaler M, Adolph TE, Ashcroft JW, *et al.* Defective ATG16L1-mediated
43 541 removal of IRE1 α drives Crohn's disease-like ileitis. *J Exp Med* 2017;**214**:401–22.
44 542 doi:10.1084/jem.20160791
45
46
47 543 25 Levine B, Kroemer G. Autophagy in the pathogenesis of disease. *Cell* 2008;**132**:27–42.
48 544 doi:10.1016/j.cell.2007.12.018
49
50
51 545 26 Hooper KM, Barlow PG, Stevens C, *et al.* Inflammatory Bowel Disease Drugs: A Focus on
52 546 Autophagy. *J Crohns Colitis* 2017;**11**:118–27. doi:10.1093/ecco-jcc/jjw127
53
54 547 27 Ke P, Shao B-Z, Xu Z-Q, *et al.* Intestinal Autophagy and Its Pharmacological Control in
55 548 Inflammatory Bowel Disease. *Front Immunol* 2017;**7**. doi:10.3389/fimmu.2016.00695
56
57
58
59
60
61
62
63
64
65

- 549 28 Hampe J, Franke A, Rosenstiel P, *et al.* A genome-wide association scan of
1 550 nonsynonymous SNPs identifies a susceptibility variant for Crohn disease in ATG16L1. *Nat*
2 551 *Genet* 2007;**39**:207–11. doi:10.1038/ng1954
3
4
5 552 29 Cadwell K, Patel KK, Maloney NS, *et al.* Virus-Plus-Susceptibility Gene Interaction
6 553 Determines Crohn’s Disease Gene Atg16L1 Phenotypes in Intestine. *Cell* 2010;**141**:1135–
7 554 45. doi:10.1016/j.cell.2010.05.009
8
9
10 555 30 Cadwell K, Liu J, Brown SL, *et al.* A unique role for autophagy and Atg16L1 in Paneth cells
11 556 in murine and human intestine. *Nature* 2008;**456**:259–63. doi:10.1038/nature07416
12
13 557 31 Lassen KG, Kuballa P, Conway KL, *et al.* Atg16L1 T300A variant decreases selective
14 558 autophagy resulting in altered cytokine signaling and decreased antibacterial defense.
15 559 *Proc Natl Acad Sci U A* 2014;**111**:7741–6. doi:10.1073/pnas.1407001111
16
17
18 560 32 Kuballa P, Huett A, Rioux JD, *et al.* Impaired autophagy of an intracellular pathogen
19 561 induced by a Crohn’s disease associated ATG16L1 variant. *PLoS One* 2008;**3**:e3391.
20 562 doi:10.1371/journal.pone.0003391
21
22
23 563 33 Singh SB, Davis AS, Taylor GA, *et al.* Human IRGM induces autophagy to eliminate
24 564 intracellular mycobacteria. *Science* 2006;**313**:1438–41. doi:10.1126/science.1129577
25
26
27 565 34 Brest P, Lapaquette P, Souidi M, *et al.* A synonymous variant in IRGM alters a binding site
28 566 for miR-196 and causes deregulation of IRGM-dependent xenophagy in Crohn’s disease.
29 567 *Nat Genet* 2011;**43**:242–5. doi:10.1038/ng.762
30
31
32 568 35 Gardet A, Benita Y, Li C, *et al.* LRRK2 is involved in the IFN-gamma response and host
33 569 response to pathogens. *J Immunol* 2010;**185**:5577–85. doi:10.4049/jimmunol.1000548
34
35
36 570 36 Marcuzzi A, Bianco AM, Girardelli M, *et al.* Genetic and functional profiling of Crohn’s
37 571 disease: autophagy mechanism and susceptibility to infectious diseases. *Biomed Res Int*
38 572 2013;**2013**:297501. doi:10.1155/2013/297501
39
40
41 573 37 Hugot JP, Chamaillard M, Zouali H, *et al.* Association of NOD2 leucine-rich repeat variants
42 574 with susceptibility to Crohn’s disease. *Nature* 2001;**411**:599–603. doi:10.1038/35079107
43
44 575 38 Rioux JD, Xavier RJ, Taylor KD, *et al.* Genome-wide association study identifies new
45 576 susceptibility loci for Crohn disease and implicates autophagy in disease pathogenesis.
46 577 *Nat Genet* 2007;**39**:596–604. doi:10.1038/ng2032
47
48
49 578 39 Weersma RK, Stokkers PCF, van Bodegraven AA, *et al.* Molecular prediction of disease risk
50 579 and severity in a large Dutch Crohn’s disease cohort. *Gut* 2009;**58**:388–95.
51 580 doi:10.1136/gut.2007.144865
52
53
54 581 40 Travassos LH, Carneiro LAM, Ramjeet M, *et al.* Nod1 and Nod2 direct autophagy by
55 582 recruiting ATG16L1 to the plasma membrane at the site of bacterial entry. *Nat Immunol*
56 583 2010;**11**:55–62. doi:10.1038/ni.1823
57
58
59
60
61
62
63
64
65

- 584 41 Chauhan S, Mandell MA, Deretic V. IRGM governs the core autophagy machinery to
1 585 conduct antimicrobial defense. *Mol Cell* 2015;**58**:507–21.
2 586 doi:10.1016/j.molcel.2015.03.020
3
4
5 587 42 Cooney R, Baker J, Brain O, *et al.* NOD2 stimulation induces autophagy in dendritic cells
6 588 influencing bacterial handling and antigen presentation. *Nat Med* 2010;**16**:90–7.
7 589 doi:10.1038/nm.2069
8
9
10 590 43 Homer CR, Richmond AL, Rebert NA, *et al.* ATG16L1 and NOD2 interact in an autophagy-
11 591 dependent, anti-bacterial pathway implicated in Crohn's disease pathogenesis.
12 592 *Gastroenterology* 2010;**139**:1630-1641.e2. doi:10.1053/j.gastro.2010.07.006
13
14 593 44 Wolfkamp SC, Verseyden C, Vogels EW, *et al.* ATG16L1 and NOD2 polymorphisms
15 594 enhance phagocytosis in monocytes of Crohn's disease patients., ATG16L1 and NOD2
16 595 polymorphisms enhance phagocytosis in monocytes of Crohn's disease patients. *World J*
17 596 *Gastroenterol World J Gastroenterol WJG* 2014;**20**, **20**:2664, 2664–72.
18 597 doi:10.3748/wjg.v20.i10.2664, 10.3748/wjg.v20.i10.2664
19
20
21
22 598 45 Adolph TE, Tomczak MF, Niederreiter L, *et al.* Paneth cells as a site of origin for intestinal
23 599 inflammation. *Nature* Published Online First: 2 October 2013. doi:10.1038/nature12599
24
25
26 600 46 Deuring JJ, Fuhler GM, Konstantinov SR, *et al.* Genomic ATG16L1 risk allele-restricted
27 601 Paneth cell ER stress in quiescent Crohn's disease. *Gut* 2014;**63**:1081–91.
28 602 doi:10.1136/gutjnl-2012-303527
29
30
31 603 47 Kobayashi KS. Nod2-Dependent Regulation of Innate and Adaptive Immunity in the
32 604 Intestinal Tract. *Science* 2005;**307**:731–4. doi:10.1126/science.1104911
33
34 605 48 Wehkamp J, Salzman NH, Porter E, *et al.* Reduced Paneth cell alpha-defensins in ileal
35 606 Crohn's disease. *Proc Natl Acad Sci U S A* 2005;**102**:18129–34.
36 607 doi:10.1073/pnas.0505256102
37
38
39 608 49 Liu B, Gulati AS, Cantillana V, *et al.* Irgm1-deficient mice exhibit Paneth cell abnormalities
40 609 and increased susceptibility to acute intestinal inflammation. *Am J Physiol - Gastrointest*
41 610 *Liver Physiol* 2013;**305**:G573–84. doi:10.1152/ajpgi.00071.2013
42
43
44 611 50 Zhang Q, Pan Y, Yan R, *et al.* Commensal bacteria direct selective cargo sorting to promote
45 612 symbiosis. *Nat Immunol* 2015;**16**:918–26. doi:10.1038/ni.3233
46
47
48 613 51 Saitoh T, Fujita N, Jang MH, *et al.* Loss of the autophagy protein Atg16L1 enhances
49 614 endotoxin-induced IL-1 β production. *Nature* 2008;**456**:264–8. doi:10.1038/nature07383
50
51
52 615 52 Zhang H, Zheng L, McGovern DPB, *et al.* Myeloid ATG16L1 Facilitates Host-Bacteria
53 616 Interactions in Maintaining Intestinal Homeostasis. *J Immunol Baltim Md 1950*
54 617 2017;**198**:2133–46. doi:10.4049/jimmunol.1601293
55
56
57 618 53 Mizoguchi A, Bhan AK. Immunobiology of B Cells in Inflammatory Bowel Disease. In:
58 619 *Crohn's Disease and Ulcerative Colitis*. Springer, Boston, MA 2012. 161–8.
59 620 doi:10.1007/978-1-4614-0998-4_12
60
61
62
63
64
65

- 621 54 Maeda S, Hsu L-C, Liu H, *et al.* Nod2 mutation in Crohn's disease potentiates NF-kappaB
1 622 activity and IL-1beta processing. *Science* 2005;**307**:734–8. doi:10.1126/science.1103685
2
- 3 623 55 Liu Z, Lee J, Krummey S, *et al.* The kinase LRRK2 is a regulator of the transcription factor
4 624 NFAT that modulates the severity of inflammatory bowel disease. *Nat Immunol*
5 625 2011;**12**:1063–70. doi:10.1038/ni.2113
6 626
- 7 626 56 Sano R, Reed JC. ER stress-induced cell death mechanisms. *Biochim Biophys Acta*
8 627 2013;**1833**. doi:10.1016/j.bbamcr.2013.06.028
9
- 10 628 57 Cao SS. Endoplasmic Reticulum Stress and Unfolded Protein Response in Inflammatory
11 629 Bowel Disease: *Inflamm Bowel Dis* 2015;**21**:636–44.
12 630 doi:10.1097/MIB.0000000000000238
13 631
- 14 631 58 Guan B-J, Krokowski D, Majumder M, *et al.* Translational control during endoplasmic
15 632 reticulum stress beyond phosphorylation of the translation initiation factor eIF2 α . *J Biol*
16 633 *Chem* 2014;**289**:12593–611. doi:10.1074/jbc.M113.543215
17 634
- 18 634 59 Vatter KM, Wek RC. Reinitiation involving upstream ORFs regulates ATF4 mRNA
19 635 translation in mammalian cells. *Proc Natl Acad Sci U S A* 2004;**101**:11269–74.
20 636 doi:10.1073/pnas.0400541101
21 637
- 22 637 60 Harding HP, Zhang Y, Bertolotti A, *et al.* Perk Is Essential for Translational Regulation and
23 638 Cell Survival during the Unfolded Protein Response. *Mol Cell* 2000;**5**:897–904.
24 639 doi:10.1016/S1097-2765(00)80330-5
25 640
- 26 640 61 Nishitoh H. CHOP is a multifunctional transcription factor in the ER stress response. *J*
27 641 *Biochem (Tokyo)* 2012;**151**:217–9. doi:10.1093/jb/mvr143
28 642
- 29 642 62 Wang XZ, Harding HP, Zhang Y, *et al.* Cloning of mammalian Ire1 reveals diversity in the
30 643 ER stress responses. *EMBO J* 1998;**17**:5708–17. doi:10.1093/emboj/17.19.5708
31 644
- 32 644 63 Shamu CE, Walter P. Oligomerization and phosphorylation of the Ire1p kinase during
33 645 intracellular signaling from the endoplasmic reticulum to the nucleus. *EMBO J*
34 646 1996;**15**:3028–39.
35 647
- 36 647 64 Tirasophon W, Lee K, Callaghan B, *et al.* The endoribonuclease activity of mammalian IRE1
37 648 autoregulates its mRNA and is required for the unfolded protein response. *Genes Dev*
38 649 2000;**14**:2725–36.
39 650
- 40 650 65 Calfon M, Zeng H, Urano F, *et al.* IRE1 couples endoplasmic reticulum load to secretory
41 651 capacity by processing the XBP-1 mRNA. *Nature* 2002;**415**:92–6. doi:10.1038/415092a
42 652
- 43 652 66 Lee A-H, Iwakoshi NN, Glimcher LH. XBP-1 Regulates a Subset of Endoplasmic Reticulum
44 653 Resident Chaperone Genes in the Unfolded Protein Response. *Mol Cell Biol*
45 654 2003;**23**:7448–59. doi:10.1128/MCB.23.21.7448-7459.2003
46 655
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 655 67 Lee K, Tirasophon W, Shen X, *et al.* IRE1-mediated unconventional mRNA splicing and S2P-
1 656 mediated ATF6 cleavage merge to regulate XBP1 in signaling the unfolded protein
2 657 response. *Genes Dev* 2002;**16**:452–66. doi:10.1101/gad.964702
3
4
5 658 68 Yoshida H, Matsui T, Yamamoto A, *et al.* XBP1 mRNA is induced by ATF6 and spliced by
6 659 IRE1 in response to ER stress to produce a highly active transcription factor. *Cell*
7 660 2001;**107**:881–91.
8
9
10 661 69 Hollien J. Decay of Endoplasmic Reticulum-Localized mRNAs During the Unfolded Protein
11 662 Response. *Science* 2006;**313**:104–7. doi:10.1126/science.1129631
12
13 663 70 Shen J, Chen X, Hendershot L, *et al.* ER stress regulation of ATF6 localization by
14 664 dissociation of BiP/GRP78 binding and unmasking of Golgi localization signals. *Dev Cell*
15 665 2002;**3**:99–111.
16
17
18 666 71 Haze K, Yoshida H, Yanagi H, *et al.* Mammalian transcription factor ATF6 is synthesized as
19 667 a transmembrane protein and activated by proteolysis in response to endoplasmic
20 668 reticulum stress. *Mol Biol Cell* 1999;**10**:3787–99.
21
22
23 669 72 Li M, Baumeister P, Roy B, *et al.* ATF6 as a transcription activator of the endoplasmic
24 670 reticulum stress element: thapsigargin stress-induced changes and synergistic
25 671 interactions with NF-Y and YY1. *Mol Cell Biol* 2000;**20**:5096–106.
26
27
28 672 73 Ye J, Rawson RB, Komuro R, *et al.* ER stress induces cleavage of membrane-bound ATF6
29 673 by the same proteases that process SREBPs. *Mol Cell* 2000;**6**:1355–64.
30
31
32 674 74 Hirsch I, Weiwad M, Prell E, *et al.* ERp29 deficiency affects sensitivity to apoptosis via
33 675 impairment of the ATF6-CHOP pathway of stress response. *Apoptosis Int J Program Cell*
34 676 *Death* 2014;**19**:801–15. doi:10.1007/s10495-013-0961-0
35
36
37 677 75 McGuckin MA, Eri RD, Das I, *et al.* ER stress and the unfolded protein response in intestinal
38 678 inflammation. *Am J Physiol - Gastrointest Liver Physiol* 2010;**298**:G820–32.
39 679 doi:10.1152/ajpgi.00063.2010
40
41
42 680 76 Deuring JJ, de Haar C, Koelewijn CL, *et al.* Absence of ABCG2-mediated mucosal
43 681 detoxification in patients with active inflammatory bowel disease is due to impeded
44 682 protein folding. *Biochem J* 2012;**441**:87–93. doi:10.1042/BJ20111281
45
46
47 683 77 Rolhion N, Barnich N, Bringer M-A, *et al.* Abnormally expressed ER stress response
48 684 chaperone Gp96 in CD favours adherent-invasive Escherichia coli invasion. *Gut*
49 685 2010;**59**:1355–62. doi:10.1136/gut.2010.207456
50
51
52 686 78 Shkoda A, Ruiz PA, Daniel H, *et al.* Interleukin-10 Blocked Endoplasmic Reticulum Stress
53 687 in Intestinal Epithelial Cells: Impact on Chronic Inflammation. *Gastroenterology*
54 688 2007;**132**:190–207. doi:10.1053/j.gastro.2006.10.030
55
56
57 689 79 Niederreiter L, Fritz TMJ, Adolph TE, *et al.* ER stress transcription factor Xbp1 suppresses
58 690 intestinal tumorigenesis and directs intestinal stem cells. *J Exp Med* 2013;**210**:2041–56.
59 691 doi:10.1084/jem.20122341
60
61
62
63
64
65

- 692 80 Martinon F, Chen X, Lee A-H, *et al.* TLR activation of the transcription factor XBP1
1 693 regulates innate immune responses in macrophages. *Nat Immunol* 2010;**11**:411–8.
2 694 doi:10.1038/ni.1857
3
4
5 695 81 Bertolotti A, Wang X, Novoa I, *et al.* Increased sensitivity to dextran sodium sulfate colitis
6 696 in IRE1 β -deficient mice. *J Clin Invest* 2001;**107**:585–93.
7
8
9 697 82 Tsuru A, Fujimoto N, Takahashi S, *et al.* Negative feedback by IRE1 β optimizes mucin
10 698 production in goblet cells. *Proc Natl Acad Sci U S A* 2013;**110**:2864–9.
11 699 doi:10.1073/pnas.1212484110
12
13 700 83 Cao SS, Wang M, Harrington JC, *et al.* Phosphorylation of eIF2 α Is Dispensable for
14 701 Differentiation but Required at a Posttranscriptional Level for Paneth Cell Function and
15 702 Intestinal Homeostasis in Mice: *Inflamm Bowel Dis* 2014;**20**:712–22.
16 703 doi:10.1097/MIB.0000000000000010
17
18
19 704 84 Cao SS, Zimmermann EM, Chuang B, *et al.* The Unfolded Protein Response and Chemical
20 705 Chaperones Reduce Protein Misfolding and Colitis in Mice. *Gastroenterology*
21 706 2013;**144**:989-1000.e6. doi:10.1053/j.gastro.2013.01.023
22
23
24 707 85 Brandl K, Rutschmann S, Li X, *et al.* Enhanced sensitivity to DSS colitis caused by a
25 708 hypomorphic Mbtps1 mutation disrupting the ATF6-driven unfolded protein response.
26 709 *Proc Natl Acad Sci U S A* 2009;**106**:3300–5. doi:10.1073/pnas.0813036106
27
28
29 710 86 Moehle C, Ackermann N, Langmann T, *et al.* Aberrant intestinal expression and allelic
30 711 variants of mucin genes associated with inflammatory bowel disease. *J Mol Med Berl Ger*
31 712 2006;**84**:1055–66. doi:10.1007/s00109-006-0100-2
32
33
34 713 87 Eri RD, Adams RJ, Tran TV, *et al.* An intestinal epithelial defect conferring ER stress results
35 714 in inflammation involving both innate and adaptive immunity. *Mucosal Immunol*
36 715 2011;**4**:354–64. doi:10.1038/mi.2010.74
37
38
39 716 88 Asada R, Saito A, Kawasaki N, *et al.* The Endoplasmic Reticulum Stress Transducer OASIS
40 717 Is involved in the Terminal Differentiation of Goblet Cells in the Large Intestine. *J Biol*
41 718 *Chem* 2012;**287**:8144–53. doi:10.1074/jbc.M111.332593
42
43
44 719 89 Hino K, Saito A, Asada R, *et al.* Increased Susceptibility to Dextran Sulfate Sodium-Induced
45 720 Colitis in the Endoplasmic Reticulum Stress Transducer OASIS Deficient Mice. *PLoS ONE*
46 721 2014;**9**. doi:10.1371/journal.pone.0088048
47
48
49 722 90 Zhao F, Edwards R, Dizon D, *et al.* Disruption of Paneth and goblet cell homeostasis and
50 723 increased endoplasmic reticulum stress in *Agr2* $^{-/-}$ mice. *Dev Biol* 2010;**338**:270–9.
51 724 doi:10.1016/j.ydbio.2009.12.008
52
53
54 725 91 Shimodaira Y, Takahashi S, Kinouchi Y, *et al.* Modulation of endoplasmic reticulum (ER)
55 726 stress-induced autophagy by C/EBP homologous protein (CHOP) and inositol-requiring
56 727 enzyme 1 α (IRE1 α) in human colon cancer cells. *Biochem Biophys Res Commun*
57 728 2014;**445**:524–33. doi:10.1016/j.bbrc.2014.02.054
58
59
60
61
62
63
64
65

- 729 92 Margariti A, Li H, Chen T, *et al.* XBP1 mRNA Splicing Triggers an Autophagic Response in
1 730 Endothelial Cells through BECLIN-1 Transcriptional Activation. *J Biol Chem* 2013;**288**:859–
2 731 72. doi:10.1074/jbc.M112.412783
3
4
5 732 93 Hetz C, Thielen P, Matus S, *et al.* XBP-1 deficiency in the nervous system protects against
6 733 amyotrophic lateral sclerosis by increasing autophagy. *Genes Dev* 2009;**23**:2294–306.
7 734 doi:10.1101/gad.1830709
8
9
10 735 94 Avivar-Valderas A, Bobrovnikova-Marjon E, Diehl JA, *et al.* Regulation of autophagy during
11 736 ECM detachment is linked to a selective inhibition of mTORC1 by PERK. *Oncogene*
12 737 2013;**32**:4932–40. doi:10.1038/onc.2012.512
13
14
15 738 95 Ji G, Yu N, Xue X, *et al.* PERK-mediated Autophagy in Osteosarcoma Cells Resists ER Stress-
16 739 induced Cell Apoptosis. *Int J Biol Sci* 2015;**11**:803–12. doi:10.7150/ijbs.11100
17
18 740 96 Jia X-E, Ma K, Xu T, *et al.* Mutation of kri1l causes definitive hematopoiesis failure via
19 741 PERK-dependent excessive autophagy induction. *Cell Res* 2015;**25**:946.
20
21
22 742 97 Kouroku Y, Fujita E, Tanida I, *et al.* ER stress (PERK/eIF2 [alpha] phosphorylation) mediates
23 743 the polyglutamine-induced LC3 conversion, an essential step for autophagy formation.
24 744 *Cell Death Differ* 2007;**14**:230.
25
26
27 745 98 Moon H, Kim B, Gwak H, *et al.* Autophagy and protein kinase RNA-like endoplasmic
28 746 reticulum kinase (PERK)/eukaryotic initiation factor 2 alpha kinase (eIF2 α) pathway
29 747 protect ovarian cancer cells from metformin-induced apoptosis: THE EFFECT OF
30 748 METFORMIN ON AUTOPHAGY AND PERK. *Mol Carcinog* 2016;**55**:346–56.
31 749 doi:10.1002/mc.22284
32
33
34 750 99 Zhao C, Yin S, Dong Y, *et al.* Autophagy-dependent EIF2AK3 activation compromises
35 751 ursolic acid-induced apoptosis through upregulation of MCL1 in MCF-7 human breast
36 752 cancer cells. *Autophagy* 2013;**9**:196–207. doi:10.4161/auto.22805
37
38
39 753 100 Laplante M, Sabatini DM. mTOR Signaling in Growth Control and Disease. *Cell*
40 754 2012;**149**:274–93. doi:10.1016/j.cell.2012.03.017
41
42
43 755 101 B'chir W, Maurin A-C, Carraro V, *et al.* The eIF2 α /ATF4 pathway is essential for stress-
44 756 induced autophagy gene expression. *Nucleic Acids Res* 2013;**41**:7683–99.
45 757 doi:10.1093/nar/gkt563
46
47
48 758 102 Avivar-Valderas A, Salas E, Bobrovnikova-Marjon E, *et al.* PERK Integrates Autophagy and
49 759 Oxidative Stress Responses To Promote Survival during Extracellular Matrix Detachment.
50 760 *Mol Cell Biol* 2011;**31**:3616–29. doi:10.1128/MCB.05164-11
51
52
53 761 103 Rouschop KMA, van den Beucken T, Dubois L, *et al.* The unfolded protein response
54 762 protects human tumor cells during hypoxia through regulation of the autophagy genes
55 763 MAP1LC3B and ATG5. *J Clin Invest* 2010;**120**:127–41. doi:10.1172/JCI40027
56
57
58 764 104 Rzymiski T, Milani M, Pike L, *et al.* Regulation of autophagy by ATF4 in response to severe
59 765 hypoxia. *Oncogene* 2010;**29**:4424–35. doi:10.1038/onc.2010.191
60
61
62
63
64
65

- 766 105 Gade P, Manjegowda SB, Nallar SC, *et al.* Regulation of the Death-Associated Protein
1 767 Kinase 1 Expression and Autophagy via ATF6 Requires Apoptosis Signal-Regulating Kinase
2 768 1. *Mol Cell Biol* 2014;**34**:4033–48. doi:10.1128/MCB.00397-14
3
4
5 769 106 Gade P, Ramachandran G, Maachani UB, *et al.* An IFN- γ -stimulated ATF6-C/EBP- β -
6 770 signaling pathway critical for the expression of Death Associated Protein Kinase 1 and
7 771 induction of autophagy. *Proc Natl Acad Sci U S A* 2012;**109**:10316–21.
8 772 doi:10.1073/pnas.1119273109
9
10
11 773 107 Lopes F, Keita ÅV, Saxena A, *et al.* ER-stress mobilization of death-associated protein
12 774 kinase-1-dependent xenophagy counteracts mitochondria stress-induced epithelial
13 775 barrier dysfunction. *J Biol Chem* 2018;**293**:3073–87. doi:10.1074/jbc.RA117.000809
14
15
16 776 108 Appenzeller-Herzog C, Hall MN. Bidirectional crosstalk between endoplasmic reticulum
17 777 stress and mTOR signaling. *Trends Cell Biol* 2012;**22**:274–82.
18 778 doi:10.1016/j.tcb.2012.02.006
19
20
21 779 109 Yamazaki H, Hiramatsu N, Hayakawa K, *et al.* Activation of the Akt-NF-kappaB pathway
22 780 by subtilase cytotoxin through the ATF6 branch of the unfolded protein response. *J*
23 781 *Immunol Baltim Md 1950* 2009;**183**:1480–7. doi:10.4049/jimmunol.0900017
24
25
26 782 110 Hart LS, Cunningham JT, Datta T, *et al.* ER stress-mediated autophagy promotes Myc-
27 783 dependent transformation and tumor growth. *J Clin Invest* 2012;**122**:4621–34.
28 784 doi:10.1172/JCI62973
29
30
31 785 111 Li J, Ni M, Lee B, *et al.* The unfolded protein response regulator GRP78/BiP is required
32 786 for endoplasmic reticulum integrity and stress-induced autophagy in mammalian cells.
33 787 *Cell Death Differ* 2008;**15**:1460–71. doi:10.1038/cdd.2008.81
34
35
36 788 112 Ogata M, Hino S, Saito A, *et al.* Autophagy is activated for cell survival after endoplasmic
37 789 reticulum stress. *Mol Cell Biol* 2006;**26**:9220–31. doi:10.1128/mcb.01453-06
38
39
40 790 113 Wang W, Kang H, Zhao Y, *et al.* Targeting autophagy sensitizes BRAF-mutant thyroid
41 791 cancer to vemurafenib. *J Clin Endocrinol Metab* 2016;**;**jc.2016-1999. doi:10.1210/jc.2016-
42 792 1999
43
44
45 793 114 Keestra-Gounder AM, Byndloss MX, Seyffert N, *et al.* NOD1 and NOD2 signalling links ER
46 794 stress with inflammation. *Nature* 2016;**532**:394–7. doi:10.1038/nature17631
47
48
49 795 115 Kaneko M, Niinuma Y, Nomura Y. Activation Signal of Nuclear Factor- κ B in Response to
50 796 Endoplasmic Reticulum Stress is Transduced via IRE1 and Tumor Necrosis Factor
51 797 Receptor-Associated Factor 2. *Biol Pharm Bull* 2003;**26**:931–5. doi:10.1248/bpb.26.931
52
53 798 116 Urano F, Wang X, Bertolotti A, *et al.* Coupling of Stress in the ER to Activation of JNK
54 799 Protein Kinases by Transmembrane Protein Kinase IRE1. *Science* 2000;**287**:664–6.
55 800 doi:10.1126/science.287.5453.664
56
57
58
59
60
61
62
63
64
65

- 801 117 Castillo K, Rojas-Rivera D, Lisbona F, *et al.* BAX inhibitor-1 regulates autophagy by
1 802 controlling the IRE1 α branch of the unfolded protein response. *EMBO J* 2011;**30**:4465–78.
2 803 doi:10.1038/emboj.2011.318
3
4
5 804 118 Ding W-X, Ni H-M, Gao W, *et al.* Linking of Autophagy to Ubiquitin-Proteasome System
6 805 Is Important for the Regulation of Endoplasmic Reticulum Stress and Cell Viability. *Am J*
7 806 *Pathol* 2007;**171**:513–24. doi:10.2353/ajpath.2007.070188
8
9
10 807 119 Moretti J, Roy S, Bozec D, *et al.* STING Senses Microbial Viability to Orchestrate Stress-
11 808 Mediated Autophagy of the Endoplasmic Reticulum. *Cell* Published Online First: October
12 809 2017. doi:10.1016/j.cell.2017.09.034
13
14
15 810 120 NHS CB. 2013/14 NHS Standard Contract for Colorectal: Complex Inflammatory Bowel
16 811 Disease (Adult). 2013.
17
18 812 121 Bassi A, Dodd S, Williamson P, *et al.* Cost of illness of inflammatory bowel disease in the
19 813 UK: a single centre retrospective study. *Gut* 2004;**53**:1471–8.
20 814 doi:10.1136/gut.2004.041616
21
22
23 815 122 Bernstein CN, Loftus EV Jr, Ng SC, *et al.* Hospitalisations and surgery in Crohn’s disease.
24 816 *Gut* 2012;**61**:622–9. doi:10.1136/gutjnl-2011-301397
25
26
27 817 123 Matsuda C, Ito T, Song J, *et al.* Therapeutic effect of a new immunosuppressive agent,
28 818 everolimus, on interleukin-10 gene-deficient mice with colitis. *Clin Exp Immunol*
29 819 2007;**148**:348–59. doi:10.1111/j.1365-2249.2007.03345.x
30
31
32 820 124 Massey DC, Bredin F, Parkes M. Use of sirolimus (rapamycin) to treat refractory Crohn’s
33 821 disease. *Gut* 2008;**57**:1294–6. doi:10.1136/gut.2008.157297
34
35
36 822 125 Dumortier J, Lapalus M-G, Guillaud O, *et al.* Everolimus for refractory Crohn’s disease: A
37 823 case report: *Inflamm Bowel Dis* 2008;**14**:874–7. doi:10.1002/ibd.20395
38
39 824 126 Mutalib M, Borrelli O, Blackstock S, *et al.* The use of sirolimus (rapamycin) in the
40 825 management of refractory inflammatory bowel disease in children. *J Crohns Colitis*
41 826 2014;**8**:1730–4. doi:10.1016/j.crohns.2014.08.014
42
43
44 827 127 Reinisch W, Panés J, Lémann M, *et al.* A multicenter, randomized, double-blind trial of
45 828 everolimus versus azathioprine and placebo to maintain steroid-induced remission in
46 829 patients with moderate-to-severe active Crohn’s disease. *Am J Gastroenterol*
47 830 2008;**103**:2284–92. doi:10.1111/j.1572-0241.2008.02024.x
48
49
50 831 128 Ha J, Kim J. Novel pharmacological modulators of autophagy: an updated patent review
51 832 (2012-2015). *Expert Opin Ther Pat* 2016;**26**:1273–89.
52 833 doi:10.1080/13543776.2016.1217996
53
54
55 834 129 Galluzzi L, Bravo-San Pedro JM, Levine B, *et al.* Pharmacological modulation of
56 835 autophagy: therapeutic potential and persisting obstacles. *Nat Rev Drug Discov*
57 836 2017;**16**:487–511. doi:10.1038/nrd.2017.22
58
59
60
61
62
63
64
65

- 837 130 Rauthan M, Pilon M. A chemical screen to identify inducers of the mitochondrial
1 838 unfolded protein response in *C. elegans*. *Worm* 2015;**4**.
2 839 doi:10.1080/21624054.2015.1096490
3
4
5 840 131 Boyce M, Bryant KF, Jousse C, *et al*. A selective inhibitor of eIF2alpha dephosphorylation
6 841 protects cells from ER stress. *Science* 2005;**307**:935–9. doi:10.1126/science.1101902
7
8 842 132 Okazaki T, Nishio A, Takeo M, *et al*. Inhibition of the dephosphorylation of eukaryotic
9 843 initiation factor 2 α ameliorates murine experimental colitis. *Digestion* 2014;**90**:167–78.
10 844 doi:10.1159/000366414
11
12
13 845 133 Crespo I, San-Miguel B, Prause C, *et al*. Glutamine treatment attenuates endoplasmic
14 846 reticulum stress and apoptosis in TNBS-induced colitis. *PLoS One* 2012;**7**:e50407.
15 847 doi:10.1371/journal.pone.0050407
16
17
18 848 134 Yang L, Sha H, Davisson RL, *et al*. Phenformin activates the unfolded protein response in
19 849 an AMP-activated protein kinase (AMPK)-dependent manner. *J Biol Chem*
20 850 2013;**288**:13631–8. doi:10.1074/jbc.M113.462762
21
22
23 851 135 Kim H, Moon SY, Kim J-S, *et al*. Activation of AMP-activated protein kinase inhibits ER
24 852 stress and renal fibrosis. *Am J Physiol Renal Physiol* 2015;**308**:F226–236.
25 853 doi:10.1152/ajprenal.00495.2014
26
27
28 854 136 Hwang H-J, Jung TW, Ryu JY, *et al*. Dipeptidyl peptidase-IV inhibitor (gemigliptin) inhibits
29 855 tunicamycin-induced endoplasmic reticulum stress, apoptosis and inflammation in H9c2
30 856 cardiomyocytes. *Mol Cell Endocrinol* 2014;**392**:1–7. doi:10.1016/j.mce.2014.04.017
31
32
33 857 137 Yusta B, Baggio LL, Estall JL, *et al*. GLP-1 receptor activation improves beta cell function
34 858 and survival following induction of endoplasmic reticulum stress. *Cell Metab* 2006;**4**:391–
35 859 406. doi:10.1016/j.cmet.2006.10.001
36
37
38 860 138 Engin F, Hotamisligil GS. Restoring endoplasmic reticulum function by chemical
39 861 chaperones: an emerging therapeutic approach for metabolic diseases. *Diabetes Obes*
40 862 *Metab* 2010;**12 Suppl 2**:108–15. doi:10.1111/j.1463-1326.2010.01282.x
41
42
43 863 139 Allaire JM, Crowley SM, Law HT, *et al*. The Intestinal Epithelium: Central Coordinator of
44 864 Mucosal Immunity. *Trends Immunol* 2018;**39**:677–96. doi:10.1016/j.it.2018.04.002
45
46
47 865 140 Boyapati RK, Rossi AG, Satsangi J, *et al*. Gut mucosal DAMPs in IBD: from mechanisms to
48 866 therapeutic implications. *Mucosal Immunol* 2016;**9**:567–82. doi:10.1038/mi.2016.14
49
50
51 867 141 Boyapati RK, Dorward DA, Tamborska A, *et al*. Mitochondrial DNA Is a Pro-Inflammatory
52 868 Damage-Associated Molecular Pattern Released During Active IBD. *Inflamm Bowel Dis*
53 869 Published Online First: 1 May 2018. doi:10.1093/ibd/izy095
54
55
56 870 142 Odenwald MA, Turner JR. The intestinal epithelial barrier: a therapeutic target? *Nat Rev*
57 871 *Gastroenterol Hepatol* 2017;**14**:9–21. doi:10.1038/nrgastro.2016.169
58
59
60
61
62
63
64
65

- 872 143 Nishino K, Nishida A, Inoue R, *et al.* Analysis of endoscopic brush samples identified
1 873 mucosa-associated dysbiosis in inflammatory bowel disease. *J Gastroenterol* 2018;**53**:95–
2 874 106. doi:10.1007/s00535-017-1384-4
3
4
5 875 144 Wildenberg ME, Koelink PJ, Diederens K, *et al.* The ATG16L1 risk allele associated with
6 876 Crohn's disease results in a Rac1-dependent defect in dendritic cell migration that is
7 877 corrected by thiopurines. *Mucosal Immunol* 2017;**10**:352–60. doi:10.1038/mi.2016.65
8
9
10 878 145 Wildenberg M, Levin A, Vos C, *et al.* P668 ATG16L1 genotype is associated with response
11 879 to anti-TNF in vitro. *J Crohns Colitis* 2013;**7**:S279. doi:10.1016/s1873-9946(13)60689-3
12
13 880 146 Levin AD, Koelink PJ, Bloemendaal FM, *et al.* Autophagy Contributes to the Induction of
14 881 Anti-TNF Induced Macrophages. *J Crohns Colitis* 2016;**10**:323–9. doi:10.1093/ecco-
15 882 jcc/jjv174
16
17
18 883 147 Vos ACW, Wildenberg ME, Arijs I, *et al.* Regulatory macrophages induced by infliximab
19 884 are involved in healing in vivo and in vitro. *Inflamm Bowel Dis* 2012;**18**:401–8.
20 885 doi:10.1002/ibd.21818
21
22
23 886 148 Niess JH, Klaus J, Stephani J, *et al.* NOD2 polymorphism predicts response to treatment
24 887 in Crohn's disease--first steps to a personalized therapy. *Dig Sci* 2012;**57**:879–86.
25 888 doi:10.1007/s10620-011-1977-3
26
27
28 889 149 Chouchana L, Fernández-Ramos AA, Dumont F, *et al.* Molecular insight into thiopurine
29 890 resistance: transcriptomic signature in lymphoblastoid cell lines. *Genome Med* 2015;**7**:37.
30 891 doi:10.1186/s13073-015-0150-6
31
32
33 892 150 Ventham NT, Kennedy NA, Nimmo ER, *et al.* Beyond gene discovery in inflammatory
34 893 bowel disease: the emerging role of epigenetics. *Gastroenterology* 2013;**145**:293–308.
35 894 doi:10.1053/j.gastro.2013.05.050
36
37
38 895 151 Kalla R, Ventham NT, Kennedy NA, *et al.* MicroRNAs: new players in IBD. *Gut*
39 896 2015;**64**:504–17. doi:10.1136/gutjnl-2014-307891
40
41
42 897 152 Stevens TW, Matheeuwsen M, Lönnkvist MH, *et al.* Systematic review: predictive
43 898 biomarkers of therapeutic response in inflammatory bowel disease--Personalised
44 899 medicine in its infancy. *Aliment Pharmacol Ther* Published Online First: 30 October 2018.
45 900 doi:10.1111/apt.15033
46
47
48 901 153 Pascal V, Pozuelo M, Borruel N, *et al.* A microbial signature for Crohn's disease. *Gut*
49 902 2017;;gutjnl-2016-313235. doi:10.1136/gutjnl-2016-313235
50
51 903
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Interactions between Autophagy and the Unfolded Protein
2 Response: Implications for Inflammatory Bowel Disease

3

4 Kirsty M. Hooper¹, Peter G. Barlow¹, Paul Henderson^{2, 3¶} and Craig Stevens^{1¶*}.

5

6 1. School of Applied Sciences, Edinburgh Napier University, Sighthill Campus, Sighthill Court,
7 Edinburgh, EH11 4BN.

8 2. Child Life and Health, University of Edinburgh, Edinburgh, EH9 1UW.

9 3. Department of Paediatric Gastroenterology and Nutrition, Royal Hospital for Sick Children,
10 Edinburgh, EH9 1LF.

11 ¶Joint senior authors

12

13 Short title: Autophagy and UPR in IBD

14

15 *Address for Correspondence:

16 Dr Craig Stevens

17 School of Applied Sciences, Edinburgh Napier University, Sighthill Campus, Sighthill Court,
18 Edinburgh, EH11 4BN.

19 Email: C.Stevens@napier.ac.uk

20 Tel: 0044 131 455 2930

21

22

23 Abstract

24 Inflammatory Bowel Disease (IBD), which includes Crohn's disease (CD) and ulcerative colitis,
25 is characterised by chronic inflammation of the gastrointestinal tract. Aetiology involves a
26 combination of genetic and environmental factors resulting in abnormal immune responses
27 to intestinal microbiota. Genetic studies have strongly linked genes involved in autophagy to
28 CD, and genes involved in the unfolded protein response (UPR) to IBD. The UPR is triggered
29 in response to accumulation of misfolded proteins in the endoplasmic reticulum (ER) and
30 autophagy plays a key role to relieve ER-stress and restore homeostasis. This review
31 summarises the known interactions between autophagy and the UPR and discusses the
32 impact of these converging pathways on IBD pathogenesis. With a paucity of effective long-
33 term treatments for IBD, targeting of synergistic pathways may provide novel and more
34 effective therapeutic options.

35 **Keywords:** IBD, autophagy, unfolded protein response, ER stress.

36

37 Introduction

38 Inflammatory Bowel Disease (IBD) is a group of inflammatory diseases that includes Crohn's
39 disease (CD), ulcerative colitis (UC) and IBD unclassified (IBDU). The incidence rate for IBD is
40 approximately 50-200 in 100,000 persons per year in Western countries [1] and following
41 diagnosis the natural history of the condition is characterized by periods of relapse and
42 remission, **with symptoms** commonly **including** abdominal pain, chronic diarrhoea, weight
43 loss and lethargy [2]. CD is distinguished from UC due to the presence of submucosal or
44 transmural inflammation and macroscopic changes that often occur in a non-contiguous
45 pattern anywhere within the digestive tract [1]. UC is localised to the colon and inflammation
46 is limited to the mucosa and epithelial lining of the gastrointestinal (GI) tract [2]. Patients can
47 be diagnosed with IBDU when a conclusive distinction between CD and UC cannot be made,
48 although this may well represent a distinct sub-type. At present there is no cure for IBD and
49 medications such as corticosteroids, aminosalicylates, immunomodulators and biological
50 agents are aimed at inducing and maintaining remission of disease by modifying inflammatory
51 processes [3].

52 **The aetiopathology** of IBD is multifactorial in nature, with genetic predisposition,
53 environmental triggers (e.g. smoking, appendicectomy, diet, pollution, antibiotics and stress)
54 and a **dysregulated mucosal immune response contributing to disease** [4]. **Examination of the**
55 **gut microbiome has revealed that IBD is associated with microbial dysbiosis, including an**
56 **expansion of facultative anaerobic bacteria of the family Enterobacteriaceae** [5]. **Several**
57 **potentially causative agents have been identified, most notably *Escherichia coli* strains with**
58 **an adherent and invasive phenotype (AIEC) are associated with ileal mucosa in CD** [6].
59 Genome wide association studies (GWAS) have identified 240 IBD susceptibility loci to date

60 [7], and have confirmed association with previously recognised susceptibility genes including
61 *Nucleotide-binding oligomerisation domain-containing protein 2 (NOD2)*. GWAS have also
62 identified the strong association of CD with genes involved in the autophagy pathway,
63 including *autophagy-related protein (ATG)16L1*, *Immunity-related GTPase family M protein*
64 (*IRGM*) and *leucine rich repeat kinase 2 (LRRK2)* [8]. The strong association of IBD with
65 endoplasmic reticulum (ER) stress/Unfolded protein response (UPR) genes including *x-box-*
66 *binding protein 1 (XBP1)* [9] and genes involved in intestinal barrier function such as *MUC2*
67 [10] and *Anterior gradient 2 (AGR2)* [11] have been detected by gene targeted approaches.
68 Together, these genetic studies have led to increased research exploring links between
69 autophagy and ER stress/UPR dysregulation and IBD pathogenesis.

70 Autophagy

71

72 Autophagy is an intracellular process that plays an important housekeeping role by degrading
73 excessive, damaged or aged proteins and organelles to maintain cellular homeostasis [12].
74 Basal autophagy is tightly regulated by the coordinated activity of autophagy-related (ATG)
75 proteins [13] and constitutes an important survival mechanism induced in response to
76 multiple stress conditions such as nutrient deprivation, hypoxia, DNA damage or intracellular
77 pathogens [12]. There are three main types of autophagy in mammalian cells;
78 macroautophagy (herein referred to as autophagy), microautophagy and chaperone-
79 mediated autophagy [12].

80 When autophagy is initiated a double membrane vesicle is formed (the autophagosome)
81 around the cargo to be degraded (**Figure 1**). The mature autophagosome then fuses with a
82 lysosome to form an autophagolysosome, in which lysosomal enzymes degrade the inner

83 membrane and cargo and the resulting macromolecules are released into the cytosol for
84 recycling (Figure 1).

85 Selective types of autophagy also exist, including autophagy of microorganisms (xenophagy)
86 and autophagy of the ER membrane (ER-phagy), which use specific receptors and adaptor
87 proteins to link the cargo to the autophagy machinery [14]. For example, Sequestosome
88 1/p62-like receptors (SLRs) target cytosolic pathogens and other cargo to initiate autophagy
89 [15]. SLRs function by binding to the small regulatory protein ubiquitin on the surface of cargo
90 [16–18] and subsequently associate with the autophagy machinery via a binding motif called
91 the LC3-interacting region (LIR) [19]. Adaptor proteins, such as autophagy-linked FYVE protein
92 (ALFY), can also bind ubiquitinated pathogens via p62 to promote association with the
93 autophagy machinery [20]. To date, five main types of SLR have been described;
94 sequestosome 1/p62, optineurin [18], NBR1 (Neighbor of BRCA1 gene 1) [21], NDP52 (Nuclear
95 Domain 10 Protein 52) [17] and the NDP52-like receptor calcoco3 (Calcium-binding and
96 coiled-coil domain-containing protein 3) [22], and specific cargo receptors are important for
97 distinct types of selective autophagy. For example, a recent study has shown that the non-
98 canonical cargo receptor cell-cycle progression gene 1 (CCPG1) is essential for ER-phagy [23],
99 while another study demonstrated an integral role for optineurin in the maintenance of ER
100 homeostasis by assisting the removal of hyper-activated UPR kinases [24].

101 Autophagy and CD

102 Autophagy affects many essential cellular processes and dysregulation of autophagy has been
103 linked to a multitude of human diseases [25]. Autophagy plays an important role in both
104 innate and adaptive immune signalling pathways and loss of immune regulation is a key event

105 leading to the chronic inflammation observed in CD [26]. Impaired autophagy responses have
106 been observed in a range of cell types derived from CD patients including the specialized
107 intestinal epithelial cells (IECs) Paneth cells and goblet cells, and leukocytes, such as
108 macrophages and dendritic cells [27].

109 Functional studies have linked impaired autophagy to CD-associated genetic variants in
110 *NOD2*, *ATG16L1*, *IRGM* and *LRRK2*. The single nucleotide polymorphism (SNP) in *ATG16L1*
111 causes a single amino acid change from threonine to alanine at position 300 (*T300A*) [28],
112 which is associated with Paneth cell and goblet cell dysfunction, and significantly impairs
113 autophagic clearance of pathogens [29–32]. *IRGM* is required for the initiation of xenophagy
114 and the clearance of intracellular organisms such as *Mycobacterium tuberculosis* [33] and
115 dysregulation of *IRGM* expression compromises the control of intracellular replication of CD-
116 associated adherent invasive *Escherichia coli* (AIEC) by autophagy [34]. *LRRK2* expression is
117 increased in colonic biopsy specimens from patients with CD [35] and functionally *LRRK2* can
118 enhance NFκB-dependent transcription, while small interfering RNA [siRNA] knockdown of
119 *LRRK2* interferes with bacterial killing [35].

120 *NOD2* is a member of the Nod-like receptor (NLR) family of pattern recognition receptors
121 (PRR) and recognises a component of the bacterial cell wall muramyl dipeptide (MDP) to
122 induce innate immune responses [36]. CD-associated *NOD2* SNPs (R702W, G908R and
123 L1007fs) affect the leucine rich repeat domain disrupting interaction with MDP and
124 abrogating immune responses initiated by this receptor [37]. The immunoregulatory
125 properties of *NOD2* have also been linked to autophagy, and CD susceptibility is heightened
126 when *ATG16L1* and *NOD2* variants present in combination, causing synergistic genetic
127 epistasis [38,39]. A direct functional interaction between these proteins has been

128 determined; NOD2 was shown to recruit ATG16L1 to the plasma membrane to initiate
129 autophagy at the sites of bacterial entry [40], and in a separate study IRGM was shown to
130 regulate the formation of a complex containing NOD2 and ATG16L1 that is necessary for the
131 induction of xenophagy [41]. The interaction of IRGM with NOD2 also stimulates
132 phosphorylation cascades involving AMPK, ULK1 and Beclin1 that regulate autophagy
133 initiation complexes [41]. Cells harbouring CD-associated *NOD2* variants and/or the *ATG16L1*
134 *T300A* variant exhibit a number of disrupted functions linked to autophagy including reduced
135 production of antimicrobial peptides, enhanced pro-inflammatory responses and aberrant
136 activation of adaptive immune responses [40,42–44].

137 Significantly, abnormalities in the secretory capacity of Paneth cells are observed in mice
138 deficient for ATG16L1 [30,45,46], NOD2 [47,48], IRGM [49] and LRRK2 [50] indicating that
139 autophagy plays an essential and specific role in Paneth cell function. Despite the significant
140 effects on Paneth cell function, mouse strains developed for deficiency in functional ATG16L1
141 do not exhibit spontaneous intestinal inflammation [29–31]. In contrast, a mouse strain with
142 targeted deletion of *ATG16L1* in IECs developed a spontaneous transmural ileitis similar to
143 ileal CD [24]. Furthermore, targeted deletion of *ATG16L* in haematopoietic cells can enhance
144 susceptibility to DSS-induced acute intestinal injury in mice [51] and ATG16L1 deficiency in
145 myeloid cells in a mouse strain led to disrupted macrophage function and bacterial clearance
146 [52]. Murine models with non-functional NOD2 do not develop spontaneous colitis [53],
147 however a *NOD2* mutation similar to the L1007fs mutation increased susceptibility to DSS-
148 induced colitis in mice [54]. *Irgm1*-deficient mice also exhibit abnormalities in Paneth cells,
149 accompanied by increased susceptibility to inflammation in the colon and ileum [49]. Finally,
150 *LRRK2* deficiency confers enhanced susceptibility to experimental colitis in mice, which was

151 associated with enhanced nuclear localisation of the transcription factor nuclear factor of
152 activated T cells (NFAT1), important for regulating innate immune responses [55].

153 ER-stress and UPR signalling

154 ER stress results from accumulation of unfolded and misfolded protein in the ER, and the UPR
155 is activated to resolve ER stress and restore homeostasis. The UPR inhibits protein synthesis,
156 promotes protein re-folding, and induces degradation of unfolded and misfolded proteins
157 through ER-associated protein degradation (ERAD) and autophagy (**Figure 2**). If these survival
158 mechanisms are unsuccessful, the UPR can induce apoptosis [56]. The major regulators of the
159 UPR are the ER-membrane resident proteins PERK (protein kinase RNA-like endoplasmic
160 reticulum kinase), inositol-requiring transmembrane kinase endonuclease 1 (IRE1) and
161 activated transcription factor (ATF)6. When inactive these proteins are bound to binding
162 immunoglobulin protein (BiP), also known as glucose regulated protein 78 (GRP78) [57].
163 During ER stress, BiP binds to misfolded proteins in the ER and dissociates from the ER-
164 membrane resident proteins to allow their transition to an active state [57] (**Figure 2**).

165 When active, PERK phosphorylates elongation initiation factor 2 α (EIF2 α), to inhibit general
166 protein synthesis [58] and specifically up-regulates ATF4 [59]. ATF4 in turn transcriptionally
167 up-regulates several other UPR genes including CCAAT/enhancer-binding protein (C/EBP)
168 homologous protein (CHOP) [60,61] (**Figure 2**). CHOP is also a transcription factor that
169 regulates several other UPR genes, and under conditions of prolonged ER stress can promote
170 apoptosis [60,61].

171 IRE1 exists in two forms: IRE1 α that is ubiquitously expressed and IRE1 β that is only expressed
172 in the GI tract and lung epithelial cells [62]. During ER stress, IRE1 is activated through

173 dimerization and auto-phosphorylation [63,64]. The IRE1 α RNase domain is essential for
174 creating transcriptionally activate *XBP1* messenger RNA (mRNA) via splicing, which acts as a
175 transactivator of UPR genes [65–68] (**Figure 2**) . IRE1 endoribonuclease activity also facilitates
176 degradation of specific mRNA in a process known as RIDD (regulated IRE1-dependent decay)
177 [69].

178 ATF6 translocates to the Golgi apparatus once released from its complex with BiP [70]. This
179 allows cleavage by site 1 and site 2 proteases (S1P and S2P), which releases the
180 transcriptionally active cytoplasmic domain of ATF6 (ATF6-N) that induces UPR-associated
181 genes [71–73] (**Figure 2**). Among the ATF6 upregulated genes are *CHOP* and *XBP1* [74].

182 ER-stress, UPR and intestinal inflammation

183 Genetic studies have identified several ER-stress/UPR genes associated with IBD [75].
184 Moreover, ER-stress levels are increased in ileal and colonic biopsies from CD patients, with
185 higher than normal levels of BiP, chaperone protein Gp96, and spliced *XBP1* observed [9,76–
186 78] (**Table 1**). Several studies have focused on IRE1-XBP1 signalling in murine models. In mice
187 with targeted deletion of XBP1 in intestinal epithelial cells (IECs) (*XBP1* ^{Δ IEC} mice), spontaneous
188 inflammation of the small intestine, increased susceptibility to DSS-induced colitis and
189 elevated levels of ER stress were observed [9] (**Table 1**). Furthermore, in *XBP1* ^{Δ IEC} mice
190 increased levels of apoptosis were observed along with reduced goblet cell and Paneth cell
191 numbers, leading to decreased production of host defence peptides and higher susceptibility
192 to *Listeria monocytogenes* infection [9] (**Table 1**). *XBP1* has also been shown to suppress
193 experimental colitis-associated cancer [79], and is essential for efficient TLR-mediated pro-

194 inflammatory responses to infection in macrophages [80]. These studies support *XBP1* as a
195 key component of the protective function of IECs and macrophages.

196 Although the UPR acts to maintain ER-homeostasis, hyper-activation of certain UPR
197 components can create a pro-inflammatory state. In *XBP1^{ΔIEC}* mice increased activation of
198 IRE1 α , causes hyper activation of NF κ B, and spontaneous inflammation [45] (**Table 1**). IRE1 β
199 knock-out mice have enhanced sensitivity to DSS-induced colitis [81] and exhibit goblet cell
200 abnormalities with exaggerated MUC2 accumulation (**Table 1**). In contrast, IRE1 α knock-out
201 mice have normal goblet cells [82]. In murine Paneth cells, IRE1 α and IRE1 β have distinct roles
202 with hyper activation of IRE1 α driving CD-like ileitis, and IRE1 β having a protective role [24].

203 Association of aberrant PERK-EIF2 α and ATF6 pathways with intestinal inflammation have
204 also been identified. A mouse model expressing non-phosphorylatable EIF2 α in IECs resulted
205 in functional abnormalities in Paneth cells and increased susceptibility to *Salmonella* infection
206 and DSS-induced colitis [83] (**Table 1**). ATF6 α deficient mice exhibit increased ER stress as
207 indicated by elevated levels of BiP, ATF4, CHOP and spliced *XBP1*, which result in enhanced
208 sensitivity to DSS-induced colitis [84] (**Table 1**). Additionally, hypomorphic mutation in
209 *membrane-bound transcription factor peptidase S1P-encoding gene (Mbtps1)*, which encodes
210 the S1P responsible for cleavage of ATF6, causes enhanced susceptibility to DSS-induced
211 colitis [85]. Although there is less evidence to support a role for PERK-EIF2 α and ATF6
212 pathways in IBD pathogenesis, their importance for ER stress responses in the intestinal
213 epithelium is clear.

214 **ER-stress and intestinal barrier function**

215 In the intestinal epithelium, cells that naturally secrete large amounts of protein, such as
216 Paneth cells and goblet cells, are more susceptible to ER-stress and therefore rely heavily on
217 the UPR to maintain homeostasis. MUC2 is the major component of mucin that is produced
218 in goblet cells and secreted into the intestinal lumen. *Winnie* mice are characterised by a
219 missense mutation in *MUC2*, which causes abnormalities in goblet cells, leading to aberrant
220 mucous production and spontaneous colitis, and association with *MUC2* variants has been
221 identified in IBD patients [86]. *Winnie* mice also exhibit severe ER stress in goblet cells [10],
222 which causes up to four-fold increase in activated dendritic cells in the colonic lamina propria,
223 and aberrant adaptive immune responses associated with interleukin (IL)-23/Th17 [87].
224 Goblet cell abnormalities are also apparent in mice deficient in UPR transcription factor
225 OASIS, which causes increased ER stress and susceptibility to DSS-induced colitis [88,89].
226 AGR2 is an ER resident protein highly expressed in goblet and Paneth cells and regulates the
227 formation of disulphide bonds in mature proteins. *AGR2*^{-/-} mice exhibit a decreased number
228 of goblet cells and MUC2 production, Paneth cell abnormalities, elevated ER-stress and
229 spontaneous colitis [90]. Notably, AGR2 is decreased in patients with CD and UC [11]. These
230 studies highlight the key role of intestinal secretory cells and breakdown of intestinal barrier
231 function in IBD pathogenesis.

232 Functional intersection between autophagy and the UPR

233 The UPR and autophagy are intimately linked processes. In a range of Intestinal epithelial cells,
234 chemical ER-stress inducers activate autophagy, modulated by enhanced expression of CHOP
235 and stimulation of the IRE1 α pathway [91]. In endothelial cells, IRE1 α -dependent splicing of
236 *XBP1* mRNA activated autophagy via up-regulation of *Beclin-1*, which is a major regulator of

237 the autophagy pathway [92] (**Figure 3**). Contrary to expectation, *XBP1* deletion in a familial
238 amyotrophic lateral sclerosis mouse model increased autophagy, which enhanced clearance
239 of accumulated toxic superoxide dismutase-1 (SOD1) aggregates [93]. It was suggested that
240 in this scenario, autophagy is induced in a compensatory manner due to attenuated UPR.

241 The UPR and autophagy also intersect at the PERK-EIF2 α -ATF4 pathway [94–99]. In an *in vitro*
242 model of osteosarcoma, PERK induced autophagy via mechanistic target of rapamycin
243 (mTORC1) inhibition to promote survival in response to ER stress-conferred chemoresistance
244 to apoptosis [95] (**Figure 3**). Additionally, PERK modulates autophagy via AMPK-dependent
245 inhibition of mTORC1 in response to extracellular matrix (ECM) detachment in mammary
246 epithelial cells (MECs) [94]. One of the main functional outcomes of PERK signalling is reduced
247 protein synthesis. Inhibition of mTORC1 helps to promote this effect as mTORC1 controls
248 synthesis of ~15-20% of protein within the cell [100]. Thus, via modulation of mTORC1, PERK
249 signalling achieves dual outcomes; inhibition of protein synthesis and induction of autophagy
250 to degrade misfolded proteins.

251 During amino acid deprivation, ATF4 and CHOP can bind specific C/EBP-ATF Response
252 Elements (CAREs), also known as Amino Acid Response Elements (AAREs) and CHOP-Response
253 Elements (CHOP-REs) to induce transcription of a wide range of autophagy genes [101] (**Figure**
254 **3**). In other studies, hypoxia or ECM detachment induced PERK-dependent autophagy due to
255 autophagy gene up-regulation via ATF4 and CHOP [102–104]. This up-regulation of autophagy
256 gene transcription by the UPR was shown to replenish autophagy proteins to promote survival
257 during cellular stress [103].

258 AT6 has also been implicated mechanistically in autophagy regulation. In response to cellular
259 stress, interferon (IFN)- γ activates the Ask1 (Apoptosis signal-regulating kinase 1)/MAPK
260 (Mitogen-activated protein kinase) pathway, which phosphorylates AT6 to allow its
261 proteolytic activation [105]. AT6 interaction with C/EBP- β is essential for IFN- γ -induced up-
262 regulation of *DAPK1* (*death-associated protein kinase 1*), which can subsequently stimulate
263 autophagy [106] (**Figure 3**). Mice lacking either AT6 or Ask1 are highly susceptible to bacterial
264 infection due to defective autophagy [105,106]. Furthermore, AT6 recruitment of DAPK1 in
265 response to ER stress enhanced xenophagy in human colonic biopsies and epithelial cells,
266 which was attenuated in cells harbouring the *ATG16L1 T300A* SNP [107]. Additionally,
267 activated AT6 was shown to stimulate Akt (protein kinase B), which resulted in the inhibition
268 of mTORC1 [108,109] (**Figure 3**).

269 In a recent study in MCF-7 human breast cancer cells, ER stress induced by the chemo-
270 preventative agent ursolic acid (UA) was associated with autophagy activation [99]. UA
271 induced autophagy via MAPK1/3 signalling and subsequent promotion of PERK signalling,
272 resulting in the inhibition of apoptosis. Furthermore, a study in human ovarian cancer cells
273 showed interdependent activation of autophagy and the PERK-EIF2 α UPR pathway when
274 treated with metformin, which causes energy starvation [98]. In these scenarios an
275 unconventional relationship between autophagy and ER stress was uncovered, which remains
276 to be mechanistically solved. Nonetheless, under these circumstances the interaction of the
277 UPR and autophagy pathways has pro-survival outcomes.

278 **Convergence of autophagy, ER-stress and CD**

279 In an attempt to relieve ER-stress the UPR can induce autophagy to degrade misfolded
280 proteins, protein aggregates and damaged organelles [91,110–113]. Autophagy activity is
281 increased in highly secretory Paneth cells [45] to counterbalance high levels of ER-stress
282 [112], thus ER-stress is a significant risk in these cells when the UPR or autophagy is not
283 functional. Consistent with this, in Paneth cells of CD patients harbouring *ATG16L1 T300A* risk
284 alleles, BiP and pEIF2 α are highly expressed [46] (**Table 1**). Significantly, *ATG16L1;XBP1 ^{Δ IEC}*
285 mice develop similar phenotypic ileitis to *ATG16L1 ^{Δ IEC}* mice, but earlier in life due to increased
286 ER stress [24,45].

287 ERAD can regulate the degradation of IRE1 α to prevent accumulation of toxic IRE1 α
288 aggregates, however persistent ER-stress will inhibit ERAD degradation of IRE1 α [24]. When
289 this occurs, autophagy plays an important role in the clearance of supramolecular clusters of
290 IRE1 α (**Figure 3**). In *ATG16L1 ^{Δ IEC}* mice, development of spontaneous CD-like ileitis is associated
291 with defective autophagy resulting in toxic accumulation of IRE1 α in Paneth cells [24] (**Table**
292 **1**). Furthermore, the selective autophagy receptor optineurin interacts with IRE1 α , and
293 optineurin deficiency amplified the accumulation of IRE1 α [24]. In humans homozygous for
294 *ATG16L1 T300A*, a similar accumulation of IRE1 α was observed in intestinal epithelial crypts
295 [24] (**Table 1**). This has led to suggestion that the *ATG16L1 T300A* SNP may define a specific
296 subtype of patients with CD, characterised by Paneth cell ER-stress [46]. This synergistic and
297 compensatory relationship between the UPR and autophagy is affirmed by the presence of
298 CD-associated SNPs in *ATG16L1* and *XBP1*.

299 A recent study has demonstrated a direct link between NOD1/2 and the IRE1 α pathway in the
300 context of ER-stress-induced inflammation [114]. When active, IRE1 α stimulates the c-Jun N-
301 terminal kinase (JNK) pathway and recruits TRAF2 (TNF receptor-associated factor 2) to the

302 ER membrane to trigger NF κ B signalling [115,116] and autophagy induction [112,117,118]
303 (**Figure 3**). In mouse and human cells, ER-stress induced by chemicals or infection with
304 *Brucella abortus* and *Chlamydia muridarum* increased inflammation and IL-6 production
305 [114]. This response was dependent on NOD1/2 and receptor-interacting serine/threonine-
306 protein kinase 2 (RIPK2), but also on IRE1 α kinase activity and TRAF2-induced NF κ B signalling
307 [114]. This suggests there is a functional intersection between the IRE1 α pathway and
308 NOD1/2 signalling, which is facilitated by TRAF2 (**Figure 3**).

309 Interestingly, an additional study has shown that ER-stress responses can be modulated by
310 another innate immune sensor called stimulator of interferon genes (STING) in response to
311 cyclic-di-AMP (c-di-AMP), a vita-PAMP (pathogen associated molecular pattern) present in
312 live Gram-positive bacteria [119]. This process induces autophagy via inhibition of the major
313 autophagy suppressor mTORC1 and localisation of STING to autophagosomes.

314 Pharmacological induction of autophagy and the UPR

315 A recent review estimated IBD treatment costs of £720 million (**\$940m**) per year in the United
316 **Kingdom alone** [120], with roughly a quarter of these costs directly attributed to drug
317 treatments [121]. The efficacy of these drugs continues to come under scrutiny as response
318 to treatment often diminishes over time, with a review of worldwide cohorts estimating that
319 between 10–35% of CD patients required surgery within a year of diagnosis and up to 61% by
320 10 years [122]. In order to improve the efficacy of IBD treatment, optimization of existing
321 clinical therapies and the development of novel therapeutics is required.

322 The convergence between autophagy and UPR pathways provides new opportunity for the
323 treatment of IBD and the modulation of the UPR in combination with autophagy inducers is a

324 promising therapeutic strategy. There is evidence that inducing autophagy can have
325 therapeutic benefits for the treatment of IBD [26] with several studies investigating the utility
326 of autophagy inducers as adjuvant therapies. Rapamycin analogues, sirolimus and everolimus,
327 inhibit mTORC1 to induce autophagy and are already approved for clinical use for post-
328 transplantation (e.g. liver and renal) management. In IL-10-deficient mice, everolimus
329 treatment alleviated spontaneous colitis and reduced CD4⁺ T cells and IFN- γ [123]. In a case
330 study sirolimus improved symptoms and intestinal healing in a patient with severe refractory
331 CD [124]. In another case study, symptoms were controlled for 18 months with everolimus
332 treatment in a refractory UC patient [125]. Moreover, in a study of refractory paediatric IBD,
333 sirolimus induced clinical remission in 45% of UC patients and 100% of CD patients; albeit the
334 sample size was small [126]. Significantly, everolimus had comparable safety and tolerability
335 as azathioprine when used to maintain steroid-induced remission in a cohort of adult CD
336 patients [127]. As these mTORC1 inhibitors are already approved for clinical use, they have
337 been investigated the most extensively, however there are a plethora of novel autophagy
338 modulators that are currently being developed, characterised and patented for therapeutic
339 use in a range of diseases including IBD [128,129].

340 Recent progress has also been made to identify specific chemical inducers of the UPR. A
341 screen of 1,200 FDA-approved compounds carried out in *C.elegans* identified eight
342 compounds that induced UPR responses, four of which specifically increased mitochondrial
343 UPR [130]. The identified drugs included antirheumatic agents, antianginal calcium channel
344 blockers; androgen receptor inhibitors used for cancer therapy and tetracycline antibiotics.

345 A well-characterised modulator of the UPR, tauroursodeoxycholic acid (TUDCA), that
346 promotes protein refolding to reduce ER-stress, was shown to ameliorate DSS-induced colitis

347 in mice by decreasing ER-stress in IECs [84]. Furthermore, a selective inhibitor of eIF2 α
348 dephosphorylation protects cells from ER-stress and ameliorates murine experimental colitis
349 [131,132]. Supplementation with glutamine has also been suggested for the improvement of
350 IBD treatment, as this amino acid was shown to dampen experimental colitis in rats by
351 inhibiting ER-stress in colonic epithelial cells [133].

352 Drugs used to treat metabolic disorders have also been investigated for UPR inducing
353 properties. The biguanides metformin and phenformin have been implicated in induction of
354 the UPR and resolution of ER-stress via activation of AMPK, which subsequently stimulated
355 IRE1 α and PERK pathways [98,134,135]. Inhibitors of dipeptidyl peptidase IV (DPP4), including
356 gemigliptin, also prevented ER-stress-mediated apoptosis by promoting IRE1 α and PERK
357 pathways [136]. Furthermore, agonists of the glucagon-like peptide-1 receptor, such as
358 exenatide, relieved ER stress via up-regulation of *ATF4* expression [137]. Exogenous chemical
359 chaperones have also been explored as a method to relieve ER stress by mimicking ER
360 chaperones to promote protein transport and re-folding capacity [138].

361 Although several studies have demonstrated beneficial effects of enhancing UPR function for
362 intestinal homeostasis, future investigations should proceed with caution. For example,
363 hyper-activation of the UPR kinase IRE1 α can exacerbate intestinal inflammation, as seen in
364 patients with *ATG16L1* and *NOD2* mutations, therefore, in certain circumstances
365 pharmacological inhibition of UPR receptors would be a more effective strategy [24,45,114]

366 Of particular interest, the selective autophagy cargo receptor optineurin forms a critical link
367 between ER-stress resolution and autophagy due to its role in the degradation of IRE1 α
368 aggregates [24], and another recently identified autophagy cargo receptor that is integral for
369 resolution of ER-stress, CCPG1, mediates ER-phagy to remove damaged ER membranes [23].
370 Understanding the biology and functions of adaptors such as optineurin and CCPG1 may
371 identify novel druggable targets and expedite development of the next generation of
372 therapeutics aimed at modulation of the UPR in combination with autophagy.

373 Discussion

374 The complexity of IBD is evident from the large number of risk loci identified by genetic
375 studies, and the diverse health profile of patients that are affected. Mouse models of IBD
376 cannot emulate the human disease, however they are useful tools to explore how specific
377 gene mutations influence inflammation. Interestingly, as highlighted in **(Table 1)** the majority
378 of mouse models mimicking IBD-associated genetic risk do not develop spontaneous
379 inflammation, but rather they are sensitised to DSS-induced colitis, which acts by damaging
380 the epithelium and increasing intestinal permeability. The intestinal epithelium has
381 important immunoregulatory functions and controls the equilibrium between tolerance and
382 immunity to non-self-antigens [139]. As such breakdown of intestinal epithelial barrier
383 function and concomitant interaction with environmental factors in the lumen is a trigger for
384 inflammation. The intestinal lumen comprises a multitude of potential triggers including the
385 microbiota, dietary antigens, and luminal antigens. Additional triggers may be host-derived
386 factors that are released into the lumen as the intestinal epithelial barrier breaks down. These
387 so-called Damage-Associated Molecular Patterns (DAMPs) include intracellular proteins, such
388 as high-mobility group box 1 (HMGB1), heat-shock proteins and components derived from

389 the extracellular matrix. Examples of non-protein DAMPs include genomic DNA,
390 mitochondrial DNA, RNA, uric acid and ATP [140,141]. Not surprisingly, there is considerable
391 interest in developing novel therapeutic strategies aimed at re-establishing intestinal barrier
392 function [142] and modulation of DAMPs for the treatment of IBD [140].

393 Dysbiosis of the gut microbiome is strongly implicated in the pathogenesis of CD [143], and it
394 has been suggested that microbial dysbiosis may be an environmental trigger. A recent study
395 by Tschurtschenthaler and colleagues [24] addressed this question. Although microbial
396 dysbiosis was present in the ileum of *Atg16l1;Xbp1*^{ΔIEC} mice, such structural alteration of the
397 microbiota did not trigger ileitis but, rather, aggravated DSS-induced colitis [24]. In order to
398 understand the role of the environment in disease, determining the relative contribution of
399 genetics and a detailed characterization of environmental triggers is required.

400 Greater understanding of the genetic factors that underlie CD pathogenesis are leading to
401 improvements in treatment. Development of personalised therapies may be achieved via
402 genotyping for key SNPs in genes involved in both the autophagy and UPR pathways. IBD
403 drugs already established in the clinic have been shown to exert their effects, at least in-part,
404 through the modulation of autophagy [26] or the UPR, and establishing patient genotypes
405 may help predict response. For example, recent studies have identified an association
406 between *ATG16L1 T300A* SNP and an enhanced therapeutic effect of thiopurines [144] and
407 anti-TNF- α therapy [145]. Interestingly, the immunoregulatory effects of these drugs were
408 associated with autophagy stimulation [144,146,147] and the *T300A* genotype has been
409 associated with a subset of patients that exhibit deficiencies in both the UPR and autophagy
410 [46]. Furthermore, CD patients harbouring *NOD2* mutations associate with better clinical
411 outcomes in response to thiopurines, whereas CD patients with wild-type *NOD2* respond

412 better to steroids and anti-TNF therapy [148]. Due to the genetic complexity of IBD and
413 epistasis between genes, it is imperative that multiple genes are analysed for the purpose of
414 patient stratification. For example, a recent study identified a 32-gene transcriptomic
415 signature in lymphoblastoid cells that was able to predict lack of response to thiopurines, with
416 aberrant cell cycle control, DNA mismatched repair and RAC1-dependent mechanisms
417 implicated in thiopurine resistance [149]. Furthermore, it is increasingly clear that epigenetic,
418 microRNA and immune cell signatures among others will have a significant role to play in
419 predicting disease susceptibility and response to therapy [150–152].

420 With regards to the intestinal microbiota, a recent study has characterised microbial
421 signatures for the diagnosis of IBD that were highly sensitive and could differentiate CD
422 patients from healthy controls and UC patients. This study highlights the potential for using
423 the intestinal microbiota as a micro-biomarker [153]. Importantly, as many drugs need to be
424 metabolised and de-toxified by the gut microbiota, this approach could also have application
425 in predicting response to therapy. Given that dysregulation of autophagy and ER-stress can
426 affect the intestinal microbial environment, analysis of microbial signatures may help to
427 determine if a patient would benefit from drugs that modulate the autophagy or UPR
428 pathways.

429 To conclude, the ER-stress/UPR and autophagy pathways play a vital role in the maintenance
430 of intestinal homeostasis and breakdown of these converging pathways has been implicated
431 in persistent intestinal infections, chronic inflammation and dysregulated immune responses
432 observed in IBD. Therefore, strategies aimed at modulating these pathways simultaneously
433 may prove to be an effective therapeutic option.

434 **Funding**

435 This work was supported by a Crohn's in Childhood Research Association (CICRA) PhD
436 studentship to KMH. PH is supported by a NHS Research Scotland Career Researcher
437 Fellowship.

438 Figure Legends

439 Figure 1: Autophagy pathway and autophagosome biogenesis

440 During the initial stages of autophagy, the isolation membrane forms a double membrane
441 vesicle (the autophagosome) around the cargo to be degraded. ULK complex (ULK1-ULK2-
442 ATG13-FIP200-ATG101) and Beclin 1 (Vps34-Vps150-Beclin1) complex, through interaction
443 with ATG14, recruit autophagy proteins and complexes to the autophagosome membrane.
444 ATG12 is conjugated to ATG5 and forms a complex with ATG16L1 (ATG16L1 complex). The
445 ATG16L1 complex is proposed to specify the site of LC3 lipidation for autophagosome
446 formation. LC3 is conjugated to PE to form lipidated LC3-II and is associated with the
447 autophagosome outer membrane. Upon autophagosome closure, LC3 localises to the inner
448 membrane and other autophagy proteins and complexes dissociate for recycling. The mature
449 autophagosome then fuses with a lysosome to form an autophagolysosome, in which cargo
450 are degraded by lysosomal enzymes and subunits are recycled.

451 Figure 2: The unfolded protein response

452 BiP chaperone protein binds unfolded/misfolded proteins in the ER and dissociates from
453 transmembrane receptors upon accumulation of the toxic proteins. The transmembrane
454 receptors PERK, IRE1 α and ATF6 become activated. PERK phosphorylates EIF2 α , which
455 downregulates global translation but specifically upregulates ATF4 and CHOP that upregulate
456 UPR-associated genes. IRE1 α splices XBP1 to its active form and ATF6 is cleaved by S1P and
457 S2P to active ATF6-N, which both translocate to the nucleus to upregulate UPR-associated
458 genes. The main function of these UPR-associated genes is to increase protein refolding,

459 inhibit synthesis of new protein and degrade unfolded/misfolded proteins through autophagy
460 and ERAD.


461 **Figure 3: Intersection between autophagy and the unfolded protein response**

462 ER stress activates transmembrane receptors PERK, IRE1 α and ATF6. PERK phosphorylates
463 EIF2 α , which specifically upregulates ATF4 and CHOP that bind AAREs and CHOP-Res to
464 upregulate autophagy genes. PERK also induces autophagy via mTORC1 inhibition. IRE1 α
465 splices XBP1 to its active form, which up-regulates *Beclin-1*. IRE1 α endonuclease activity
466 activates the JNK pathway, which induces autophagy via TRAF2, NOD2 and NF κ B. Enhanced
467 autophagy degrades accumulated IRE1 α clusters. Active ATF6-N induces autophagy via
468 mTORC1 inhibition and binds C/EBP- β to up-regulate *DAPK1*.

469 **Table 1: Murine models of intestinal inflammation**


470 **Links between autophagy, ER-stress/UPR and experimental colitis/intestinal inflammation**
471 **and IBD.**

472




473

474 Figure 1



475
476 Figure 2



477
478

Figure 3

Autophagy/ UPR pathway	Murine models of intestinal inflammation	IBD patients
ATG16L1	<ul style="list-style-type: none"> • ATG16L1 deficiency caused enhanced susceptibility to experimental colitis, Paneth cell and Goblet cell dysfunction, disrupted macrophage function and significantly impairs xenophagy [29-32, 51, 52] • ATG16L1 deletion in IECs induced spontaneous transmural ileitis [24] 	<i>ATG16L1 T300A</i> CD-associated SNP [28]
NOD2	NOD2 mutation causes enhanced susceptibility to DSS-induced colitis [54] and causes Paneth cell dysfunction [47, 48]	<i>NOD2</i> CD-associated SNPs (R702W, G908R and L1007fs) [37]
IRGM	<i>Irgm1</i> deficiency causes abnormalities in Paneth cells and increased susceptibility to inflammation in the colon and ileum [49]	<i>IRGM</i> CD-associated SNP [8]
LRRK4	LRRK2 deficiency confers enhanced susceptibility to experimental colitis in mice [55] and Paneth cell abnormalities [50]	<i>LRRK4</i> CD-associated SNP [8]
IRE1 α -XBP1	<ul style="list-style-type: none"> • <i>XBP1</i> deletion causes spontaneous intestinal inflammation, abnormal Paneth and goblet cell function and increased infection [9] • <i>XBP1</i> deletion causes overactivation of IRE1α and NFκB [45] • <i>ATG16L1</i> deletion causes accumulation of IRE1α in Paneth cells resulting in CD-like ileitis [24] 	<ul style="list-style-type: none"> • <i>XBP1</i> CD-associated SNP [9] • Increased levels of spliced <i>XBP1</i>, BiP and Gp96 in CD [9, 76-78] • <i>T300A</i> SNP causes accumulation of IRE1α in intestinal crypts [24]
IRE1 β	IRE1 β deletion causes enhanced sensitivity to DSS-colitis [81], goblet cell abnormalities and MUC2 accumulation [24]	
PERK-EIF2 α	Non-phosphorylatable EIF2 α caused Paneth cell abnormalities, enhanced DSS-colitis susceptibility and increased <i>Salmonella</i> infection [83]	Increased p-EIF2 α and BiP in CD patients with <i>T300A</i> SNP [46]
ATF6	<ul style="list-style-type: none"> • ATF6 deletion enhanced DSS-colitis susceptibility [84] • Mutation in <i>Mbtps1</i> (encodes S1P) causes enhanced DSS-colitis susceptibility [85] 	
AGR2	AGR2 deletion causes decreased Goblet cells and MUC2 production, Paneth cell abnormalities, elevated ER-stress and spontaneous colitis [90]	<ul style="list-style-type: none"> • <i>AGR2</i> CD-associated SNP [11] • <i>AGR2</i> decreased in IBD [11]

480 Table 1

481 References

- 482 1 Gasparetto M, Guariso G. Highlights in IBD Epidemiology and Its Natural History in the
483 Paediatric Age. *Gastroenterol Res Pr* 2013;**2013**:829040. doi:10.1155/2013/829040
- 484 2 Fakhoury M, Negrulj R, Mooranian A, *et al.* Inflammatory bowel disease: clinical aspects
485 and treatments. *J Inflamm Res* 2014;**7**:113–20. doi:10.2147/JIR.S65979
- 486 3 Neurath MF. Current and emerging therapeutic targets for IBD. *Nat Rev Gastroenterol*
487 *Hepatol* 2017;**14**:nrgastro.2016.208. doi:10.1038/nrgastro.2016.208
- 488 4 Boyapati R, Satsangi J, Ho GT. Pathogenesis of Crohn's disease. *F1000Prime Rep*
489 2015;**7**:44. doi:10.12703/p7-44
- 490 5 Zuo T, Ng SC. The Gut Microbiota in the Pathogenesis and Therapeutics of Inflammatory
491 Bowel Disease. *Front Microbiol* 2018;**9**:2247. doi:10.3389/fmicb.2018.02247
- 492 6 Darfeuille-Michaud A, Boudeau J, Bulois P, *et al.* High prevalence of adherent-invasive
493 *Escherichia coli* associated with ileal mucosa in Crohn's disease. *Gastroenterology*
494 2004;**127**:412–21. doi:10.1053/j.gastro.2004.04.061
- 495 7 Lange KM de, Moutsianas L, Lee JC, *et al.* Genome-wide association study implicates
496 immune activation of multiple integrin genes in inflammatory bowel disease. *Nat Genet*
497 2017;**49**:256–61. doi:10.1038/ng.3760
- 498 8 Franke A, McGovern DP, Barrett JC, *et al.* Genome-wide meta-analysis increases to 71 the
499 number of confirmed Crohn's disease susceptibility loci. *Nat Genet* 2010;**42**:1118–25.
500 doi:10.1038/ng.717
- 501 9 Kaser A, Lee A-H, Franke A, *et al.* XBP1 Links ER Stress to Intestinal Inflammation and
502 Confers Genetic Risk for Human Inflammatory Bowel Disease. *Cell* 2008;**134**:743–56.
503 doi:10.1016/j.cell.2008.07.021
- 504 10 Heazlewood CK, Cook MC, Eri R, *et al.* Aberrant Mucin Assembly in Mice Causes
505 Endoplasmic Reticulum Stress and Spontaneous Inflammation Resembling Ulcerative
506 Colitis. *PLoS Med* 2008;**5**. doi:10.1371/journal.pmed.0050054
- 507 11 Zheng W, Rosenstiel P, Huse K, *et al.* Evaluation of AGR2 and AGR3 as candidate genes for
508 inflammatory bowel disease. *Genes Immun* 2006;**7**:11–8. doi:10.1038/sj.gene.6364263
- 509 12 Dikic I, Elazar Z. Mechanism and medical implications of mammalian autophagy. *Nat Rev*
510 *Mol Cell Biol* 2018;**19**:349. doi:10.1038/s41580-018-0003-4
- 511 13 Lamb CA, Yoshimori T, Tooze SA. The autophagosome: origins unknown, biogenesis
512 complex. *Nat Rev Mol Cell Biol* 2013;**14**:759–74. doi:10.1038/nrm3696
- 513 14 Zaffagnini G, Martens S. Mechanisms of Selective Autophagy. *J Mol Biol* 2016;**428**:1714–
514 24. doi:10.1016/j.jmb.2016.02.004

- 515 15 Nys K, Agostinis P, Vermeire S. Autophagy: a new target or an old strategy for the
516 treatment of Crohn's disease? *Nat Rev Gastroenterol Hepatol* 2013;**10**:395–401.
517 doi:10.1038/nrgastro.2013.66
- 518 16 Dupont N, Lacas-Gervais S, Bertout J, *et al.* Shigella phagocytic vacuolar membrane
519 remnants participate in the cellular response to pathogen invasion and are regulated by
520 autophagy. *Cell Host Microbe* 2009;**6**:137–49. doi:10.1016/j.chom.2009.07.005
- 521 17 Thurston TLM, Ryzhakov G, Bloor S, *et al.* The TBK1 adaptor and autophagy receptor
522 NDP52 restricts the proliferation of ubiquitin-coated bacteria. *Nat Immunol*
523 2009;**10**:1215–21. doi:10.1038/ni.1800
- 524 18 Wild P, Farhan H, McEwan DG, *et al.* Phosphorylation of the autophagy receptor
525 optineurin restricts Salmonella growth. *Science* 2011;**333**:228–33.
526 doi:10.1126/science.1205405
- 527 19 Johansen T, Lamark T. Selective autophagy mediated by autophagic adapter proteins.
528 *Autophagy* 2011;**7**:279–96.
- 529 20 Filimonenko M, Isakson P, Finley KD, *et al.* The selective macroautophagic degradation of
530 aggregated proteins requires the PI3P-binding protein Alfy. *Mol Cell* 2010;**38**:265–79.
531 doi:10.1016/j.molcel.2010.04.007
- 532 21 Kirkin V, Lamark T, Sou Y-S, *et al.* A role for NBR1 in autophagosomal degradation of
533 ubiquitinated substrates. *Mol Cell* 2009;**33**:505–16. doi:10.1016/j.molcel.2009.01.020
- 534 22 Newman AC, Scholefield CL, Kemp AJ, *et al.* TBK1 kinase addiction in lung cancer cells is
535 mediated via autophagy of Tax1bp1/Ndp52 and non-canonical NF- κ B signalling. *PLoS One*
536 2012;**7**:e50672. doi:10.1371/journal.pone.0050672
- 537 23 Smith MD, Harley ME, Kemp AJ, *et al.* CCPG1 Is a Non-canonical Autophagy Cargo
538 Receptor Essential for ER-Phagy and Pancreatic ER Proteostasis. *Dev Cell* 2018;**44**:217-
539 232.e11. doi:10.1016/j.devcel.2017.11.024
- 540 24 Tschurtschenthaler M, Adolph TE, Ashcroft JW, *et al.* Defective ATG16L1-mediated
541 removal of IRE1 α drives Crohn's disease-like ileitis. *J Exp Med* 2017;**214**:401–22.
542 doi:10.1084/jem.20160791
- 543 25 Levine B, Kroemer G. Autophagy in the pathogenesis of disease. *Cell* 2008;**132**:27–42.
544 doi:10.1016/j.cell.2007.12.018
- 545 26 Hooper KM, Barlow PG, Stevens C, *et al.* Inflammatory Bowel Disease Drugs: A Focus on
546 Autophagy. *J Crohns Colitis* 2017;**11**:118–27. doi:10.1093/ecco-jcc/jjw127
- 547 27 Ke P, Shao B-Z, Xu Z-Q, *et al.* Intestinal Autophagy and Its Pharmacological Control in
548 Inflammatory Bowel Disease. *Front Immunol* 2017;**7**. doi:10.3389/fimmu.2016.00695

- 549 28 Hampe J, Franke A, Rosenstiel P, *et al.* A genome-wide association scan of
550 nonsynonymous SNPs identifies a susceptibility variant for Crohn disease in ATG16L1. *Nat*
551 *Genet* 2007;**39**:207–11. doi:10.1038/ng1954
- 552 29 Cadwell K, Patel KK, Maloney NS, *et al.* Virus-Plus-Susceptibility Gene Interaction
553 Determines Crohn’s Disease Gene Atg16L1 Phenotypes in Intestine. *Cell* 2010;**141**:1135–
554 45. doi:10.1016/j.cell.2010.05.009
- 555 30 Cadwell K, Liu J, Brown SL, *et al.* A unique role for autophagy and Atg16L1 in Paneth cells
556 in murine and human intestine. *Nature* 2008;**456**:259–63. doi:10.1038/nature07416
- 557 31 Lassen KG, Kuballa P, Conway KL, *et al.* Atg16L1 T300A variant decreases selective
558 autophagy resulting in altered cytokine signaling and decreased antibacterial defense.
559 *Proc Natl Acad Sci U A* 2014;**111**:7741–6. doi:10.1073/pnas.1407001111
- 560 32 Kuballa P, Huett A, Rioux JD, *et al.* Impaired autophagy of an intracellular pathogen
561 induced by a Crohn’s disease associated ATG16L1 variant. *PLoS One* 2008;**3**:e3391.
562 doi:10.1371/journal.pone.0003391
- 563 33 Singh SB, Davis AS, Taylor GA, *et al.* Human IRGM induces autophagy to eliminate
564 intracellular mycobacteria. *Science* 2006;**313**:1438–41. doi:10.1126/science.1129577
- 565 34 Brest P, Lapaquette P, Souidi M, *et al.* A synonymous variant in IRGM alters a binding site
566 for miR-196 and causes deregulation of IRGM-dependent xenophagy in Crohn’s disease.
567 *Nat Genet* 2011;**43**:242–5. doi:10.1038/ng.762
- 568 35 Gardet A, Benita Y, Li C, *et al.* LRRK2 is involved in the IFN-gamma response and host
569 response to pathogens. *J Immunol* 2010;**185**:5577–85. doi:10.4049/jimmunol.1000548
- 570 36 Marcuzzi A, Bianco AM, Girardelli M, *et al.* Genetic and functional profiling of Crohn’s
571 disease: autophagy mechanism and susceptibility to infectious diseases. *Biomed Res Int*
572 2013;**2013**:297501. doi:10.1155/2013/297501
- 573 37 Hugot JP, Chamaillard M, Zouali H, *et al.* Association of NOD2 leucine-rich repeat variants
574 with susceptibility to Crohn’s disease. *Nature* 2001;**411**:599–603. doi:10.1038/35079107
- 575 38 Rioux JD, Xavier RJ, Taylor KD, *et al.* Genome-wide association study identifies new
576 susceptibility loci for Crohn disease and implicates autophagy in disease pathogenesis.
577 *Nat Genet* 2007;**39**:596–604. doi:10.1038/ng2032
- 578 39 Weersma RK, Stokkers PCF, van Bodegraven AA, *et al.* Molecular prediction of disease risk
579 and severity in a large Dutch Crohn’s disease cohort. *Gut* 2009;**58**:388–95.
580 doi:10.1136/gut.2007.144865
- 581 40 Travassos LH, Carneiro LAM, Ramjeet M, *et al.* Nod1 and Nod2 direct autophagy by
582 recruiting ATG16L1 to the plasma membrane at the site of bacterial entry. *Nat Immunol*
583 2010;**11**:55–62. doi:10.1038/ni.1823

- 584 41 Chauhan S, Mandell MA, Deretic V. IRGM governs the core autophagy machinery to
585 conduct antimicrobial defense. *Mol Cell* 2015;**58**:507–21.
586 doi:10.1016/j.molcel.2015.03.020
- 587 42 Cooney R, Baker J, Brain O, *et al.* NOD2 stimulation induces autophagy in dendritic cells
588 influencing bacterial handling and antigen presentation. *Nat Med* 2010;**16**:90–7.
589 doi:10.1038/nm.2069
- 590 43 Homer CR, Richmond AL, Rebert NA, *et al.* ATG16L1 and NOD2 interact in an autophagy-
591 dependent, anti-bacterial pathway implicated in Crohn's disease pathogenesis.
592 *Gastroenterology* 2010;**139**:1630-1641.e2. doi:10.1053/j.gastro.2010.07.006
- 593 44 Wolfkamp SC, Verseyden C, Vogels EW, *et al.* ATG16L1 and NOD2 polymorphisms
594 enhance phagocytosis in monocytes of Crohn's disease patients., ATG16L1 and NOD2
595 polymorphisms enhance phagocytosis in monocytes of Crohn's disease patients. *World J*
596 *Gastroenterol World J Gastroenterol WJG* 2014;**20**, **20**:2664, 2664–72.
597 doi:10.3748/wjg.v20.i10.2664, 10.3748/wjg.v20.i10.2664
- 598 45 Adolph TE, Tomczak MF, Niederreiter L, *et al.* Paneth cells as a site of origin for intestinal
599 inflammation. *Nature* Published Online First: 2 October 2013. doi:10.1038/nature12599
- 600 46 Deuring JJ, Fuhler GM, Konstantinov SR, *et al.* Genomic ATG16L1 risk allele-restricted
601 Paneth cell ER stress in quiescent Crohn's disease. *Gut* 2014;**63**:1081–91.
602 doi:10.1136/gutjnl-2012-303527
- 603 47 Kobayashi KS. Nod2-Dependent Regulation of Innate and Adaptive Immunity in the
604 Intestinal Tract. *Science* 2005;**307**:731–4. doi:10.1126/science.1104911
- 605 48 Wehkamp J, Salzman NH, Porter E, *et al.* Reduced Paneth cell alpha-defensins in ileal
606 Crohn's disease. *Proc Natl Acad Sci U S A* 2005;**102**:18129–34.
607 doi:10.1073/pnas.0505256102
- 608 49 Liu B, Gulati AS, Cantillana V, *et al.* Irgm1-deficient mice exhibit Paneth cell abnormalities
609 and increased susceptibility to acute intestinal inflammation. *Am J Physiol - Gastrointest*
610 *Liver Physiol* 2013;**305**:G573–84. doi:10.1152/ajpgi.00071.2013
- 611 50 Zhang Q, Pan Y, Yan R, *et al.* Commensal bacteria direct selective cargo sorting to promote
612 symbiosis. *Nat Immunol* 2015;**16**:918–26. doi:10.1038/ni.3233
- 613 51 Saitoh T, Fujita N, Jang MH, *et al.* Loss of the autophagy protein Atg16L1 enhances
614 endotoxin-induced IL-1 β production. *Nature* 2008;**456**:264–8. doi:10.1038/nature07383
- 615 52 Zhang H, Zheng L, McGovern DPB, *et al.* Myeloid ATG16L1 Facilitates Host-Bacteria
616 Interactions in Maintaining Intestinal Homeostasis. *J Immunol Baltim Md 1950*
617 **2017**;**198**:2133–46. doi:10.4049/jimmunol.1601293
- 618 53 Mizoguchi A, Bhan AK. Immunobiology of B Cells in Inflammatory Bowel Disease. In:
619 *Crohn's Disease and Ulcerative Colitis*. Springer, Boston, MA 2012. 161–8.
620 doi:10.1007/978-1-4614-0998-4_12

- 621 54 Maeda S, Hsu L-C, Liu H, *et al.* Nod2 mutation in Crohn's disease potentiates NF-kappaB
622 activity and IL-1beta processing. *Science* 2005;**307**:734–8. doi:10.1126/science.1103685
- 623 55 Liu Z, Lee J, Krummey S, *et al.* The kinase LRRK2 is a regulator of the transcription factor
624 NFAT that modulates the severity of inflammatory bowel disease. *Nat Immunol*
625 2011;**12**:1063–70. doi:10.1038/ni.2113
- 626 56 Sano R, Reed JC. ER stress-induced cell death mechanisms. *Biochim Biophys Acta*
627 2013;**1833**. doi:10.1016/j.bbamcr.2013.06.028
- 628 57 Cao SS. Endoplasmic Reticulum Stress and Unfolded Protein Response in Inflammatory
629 Bowel Disease: *Inflamm Bowel Dis* 2015;**21**:636–44.
630 doi:10.1097/MIB.0000000000000238
- 631 58 Guan B-J, Krokowski D, Majumder M, *et al.* Translational control during endoplasmic
632 reticulum stress beyond phosphorylation of the translation initiation factor eIF2 α . *J Biol*
633 *Chem* 2014;**289**:12593–611. doi:10.1074/jbc.M113.543215
- 634 59 Vatter KM, Wek RC. Reinitiation involving upstream ORFs regulates ATF4 mRNA
635 translation in mammalian cells. *Proc Natl Acad Sci U S A* 2004;**101**:11269–74.
636 doi:10.1073/pnas.0400541101
- 637 60 Harding HP, Zhang Y, Bertolotti A, *et al.* Perk Is Essential for Translational Regulation and
638 Cell Survival during the Unfolded Protein Response. *Mol Cell* 2000;**5**:897–904.
639 doi:10.1016/S1097-2765(00)80330-5
- 640 61 Nishitoh H. CHOP is a multifunctional transcription factor in the ER stress response. *J*
641 *Biochem (Tokyo)* 2012;**151**:217–9. doi:10.1093/jb/mvr143
- 642 62 Wang XZ, Harding HP, Zhang Y, *et al.* Cloning of mammalian Ire1 reveals diversity in the
643 ER stress responses. *EMBO J* 1998;**17**:5708–17. doi:10.1093/emboj/17.19.5708
- 644 63 Shamu CE, Walter P. Oligomerization and phosphorylation of the Ire1p kinase during
645 intracellular signaling from the endoplasmic reticulum to the nucleus. *EMBO J*
646 1996;**15**:3028–39.
- 647 64 Tirasophon W, Lee K, Callaghan B, *et al.* The endoribonuclease activity of mammalian IRE1
648 autoregulates its mRNA and is required for the unfolded protein response. *Genes Dev*
649 2000;**14**:2725–36.
- 650 65 Calfon M, Zeng H, Urano F, *et al.* IRE1 couples endoplasmic reticulum load to secretory
651 capacity by processing the XBP-1 mRNA. *Nature* 2002;**415**:92–6. doi:10.1038/415092a
- 652 66 Lee A-H, Iwakoshi NN, Glimcher LH. XBP-1 Regulates a Subset of Endoplasmic Reticulum
653 Resident Chaperone Genes in the Unfolded Protein Response. *Mol Cell Biol*
654 2003;**23**:7448–59. doi:10.1128/MCB.23.21.7448-7459.2003

- 655 67 Lee K, Tirasophon W, Shen X, *et al.* IRE1-mediated unconventional mRNA splicing and S2P-
656 mediated ATF6 cleavage merge to regulate XBP1 in signaling the unfolded protein
657 response. *Genes Dev* 2002;**16**:452–66. doi:10.1101/gad.964702
- 658 68 Yoshida H, Matsui T, Yamamoto A, *et al.* XBP1 mRNA is induced by ATF6 and spliced by
659 IRE1 in response to ER stress to produce a highly active transcription factor. *Cell*
660 2001;**107**:881–91.
- 661 69 Hollien J. Decay of Endoplasmic Reticulum-Localized mRNAs During the Unfolded Protein
662 Response. *Science* 2006;**313**:104–7. doi:10.1126/science.1129631
- 663 70 Shen J, Chen X, Hendershot L, *et al.* ER stress regulation of ATF6 localization by
664 dissociation of BiP/GRP78 binding and unmasking of Golgi localization signals. *Dev Cell*
665 2002;**3**:99–111.
- 666 71 Haze K, Yoshida H, Yanagi H, *et al.* Mammalian transcription factor ATF6 is synthesized as
667 a transmembrane protein and activated by proteolysis in response to endoplasmic
668 reticulum stress. *Mol Biol Cell* 1999;**10**:3787–99.
- 669 72 Li M, Baumeister P, Roy B, *et al.* ATF6 as a transcription activator of the endoplasmic
670 reticulum stress element: thapsigargin stress-induced changes and synergistic
671 interactions with NF-Y and YY1. *Mol Cell Biol* 2000;**20**:5096–106.
- 672 73 Ye J, Rawson RB, Komuro R, *et al.* ER stress induces cleavage of membrane-bound ATF6
673 by the same proteases that process SREBPs. *Mol Cell* 2000;**6**:1355–64.
- 674 74 Hirsch I, Weiwad M, Prell E, *et al.* ERp29 deficiency affects sensitivity to apoptosis via
675 impairment of the ATF6-CHOP pathway of stress response. *Apoptosis Int J Program Cell*
676 *Death* 2014;**19**:801–15. doi:10.1007/s10495-013-0961-0
- 677 75 McGuckin MA, Eri RD, Das I, *et al.* ER stress and the unfolded protein response in intestinal
678 inflammation. *Am J Physiol - Gastrointest Liver Physiol* 2010;**298**:G820–32.
679 doi:10.1152/ajpgi.00063.2010
- 680 76 Deuring JJ, de Haar C, Koelewijn CL, *et al.* Absence of ABCG2-mediated mucosal
681 detoxification in patients with active inflammatory bowel disease is due to impeded
682 protein folding. *Biochem J* 2012;**441**:87–93. doi:10.1042/BJ20111281
- 683 77 Rolhion N, Barnich N, Bringer M-A, *et al.* Abnormally expressed ER stress response
684 chaperone Gp96 in CD favours adherent-invasive Escherichia coli invasion. *Gut*
685 2010;**59**:1355–62. doi:10.1136/gut.2010.207456
- 686 78 Shkoda A, Ruiz PA, Daniel H, *et al.* Interleukin-10 Blocked Endoplasmic Reticulum Stress
687 in Intestinal Epithelial Cells: Impact on Chronic Inflammation. *Gastroenterology*
688 2007;**132**:190–207. doi:10.1053/j.gastro.2006.10.030
- 689 79 Niederreiter L, Fritz TMJ, Adolph TE, *et al.* ER stress transcription factor Xbp1 suppresses
690 intestinal tumorigenesis and directs intestinal stem cells. *J Exp Med* 2013;**210**:2041–56.
691 doi:10.1084/jem.20122341

- 692 80 Martinon F, Chen X, Lee A-H, *et al.* TLR activation of the transcription factor XBP1
693 regulates innate immune responses in macrophages. *Nat Immunol* 2010;**11**:411–8.
694 doi:10.1038/ni.1857
- 695 81 Bertolotti A, Wang X, Novoa I, *et al.* Increased sensitivity to dextran sodium sulfate colitis
696 in IRE1 β -deficient mice. *J Clin Invest* 2001;**107**:585–93.
- 697 82 Tsuru A, Fujimoto N, Takahashi S, *et al.* Negative feedback by IRE1 β optimizes mucin
698 production in goblet cells. *Proc Natl Acad Sci U S A* 2013;**110**:2864–9.
699 doi:10.1073/pnas.1212484110
- 700 83 Cao SS, Wang M, Harrington JC, *et al.* Phosphorylation of eIF2 α Is Dispensable for
701 Differentiation but Required at a Posttranscriptional Level for Paneth Cell Function and
702 Intestinal Homeostasis in Mice: *Inflamm Bowel Dis* 2014;**20**:712–22.
703 doi:10.1097/MIB.0000000000000010
- 704 84 Cao SS, Zimmermann EM, Chuang B, *et al.* The Unfolded Protein Response and Chemical
705 Chaperones Reduce Protein Misfolding and Colitis in Mice. *Gastroenterology*
706 2013;**144**:989-1000.e6. doi:10.1053/j.gastro.2013.01.023
- 707 85 Brandl K, Rutschmann S, Li X, *et al.* Enhanced sensitivity to DSS colitis caused by a
708 hypomorphic Mbtps1 mutation disrupting the ATF6-driven unfolded protein response.
709 *Proc Natl Acad Sci U S A* 2009;**106**:3300–5. doi:10.1073/pnas.0813036106
- 710 86 Moehle C, Ackermann N, Langmann T, *et al.* Aberrant intestinal expression and allelic
711 variants of mucin genes associated with inflammatory bowel disease. *J Mol Med Berl Ger*
712 2006;**84**:1055–66. doi:10.1007/s00109-006-0100-2
- 713 87 Eri RD, Adams RJ, Tran TV, *et al.* An intestinal epithelial defect conferring ER stress results
714 in inflammation involving both innate and adaptive immunity. *Mucosal Immunol*
715 2011;**4**:354–64. doi:10.1038/mi.2010.74
- 716 88 Asada R, Saito A, Kawasaki N, *et al.* The Endoplasmic Reticulum Stress Transducer OASIS
717 Is Involved in the Terminal Differentiation of Goblet Cells in the Large Intestine. *J Biol*
718 *Chem* 2012;**287**:8144–53. doi:10.1074/jbc.M111.332593
- 719 89 Hino K, Saito A, Asada R, *et al.* Increased Susceptibility to Dextran Sulfate Sodium-Induced
720 Colitis in the Endoplasmic Reticulum Stress Transducer OASIS Deficient Mice. *PLoS ONE*
721 2014;**9**. doi:10.1371/journal.pone.0088048
- 722 90 Zhao F, Edwards R, Dizon D, *et al.* Disruption of Paneth and goblet cell homeostasis and
723 increased endoplasmic reticulum stress in *Agr2* $^{-/-}$ mice. *Dev Biol* 2010;**338**:270–9.
724 doi:10.1016/j.ydbio.2009.12.008
- 725 91 Shimodaira Y, Takahashi S, Kinouchi Y, *et al.* Modulation of endoplasmic reticulum (ER)
726 stress-induced autophagy by C/EBP homologous protein (CHOP) and inositol-requiring
727 enzyme 1 α (IRE1 α) in human colon cancer cells. *Biochem Biophys Res Commun*
728 2014;**445**:524–33. doi:10.1016/j.bbrc.2014.02.054

- 729 92 Margariti A, Li H, Chen T, *et al.* XBP1 mRNA Splicing Triggers an Autophagic Response in
730 Endothelial Cells through BECLIN-1 Transcriptional Activation. *J Biol Chem* 2013;**288**:859–
731 72. doi:10.1074/jbc.M112.412783
- 732 93 Hetz C, Thielen P, Matus S, *et al.* XBP-1 deficiency in the nervous system protects against
733 amyotrophic lateral sclerosis by increasing autophagy. *Genes Dev* 2009;**23**:2294–306.
734 doi:10.1101/gad.1830709
- 735 94 Avivar-Valderas A, Bobrovnikova-Marjon E, Diehl JA, *et al.* Regulation of autophagy during
736 ECM detachment is linked to a selective inhibition of mTORC1 by PERK. *Oncogene*
737 2013;**32**:4932–40. doi:10.1038/onc.2012.512
- 738 95 Ji G, Yu N, Xue X, *et al.* PERK-mediated Autophagy in Osteosarcoma Cells Resists ER Stress-
739 induced Cell Apoptosis. *Int J Biol Sci* 2015;**11**:803–12. doi:10.7150/ijbs.11100
- 740 96 Jia X-E, Ma K, Xu T, *et al.* Mutation of kri1l causes definitive hematopoiesis failure via
741 PERK-dependent excessive autophagy induction. *Cell Res* 2015;**25**:946.
- 742 97 Kouroku Y, Fujita E, Tanida I, *et al.* ER stress (PERK/eIF2 [alpha] phosphorylation) mediates
743 the polyglutamine-induced LC3 conversion, an essential step for autophagy formation.
744 *Cell Death Differ* 2007;**14**:230.
- 745 98 Moon H, Kim B, Gwak H, *et al.* Autophagy and protein kinase RNA-like endoplasmic
746 reticulum kinase (PERK)/eukaryotic initiation factor 2 alpha kinase (eIF2 α) pathway
747 protect ovarian cancer cells from metformin-induced apoptosis: THE EFFECT OF
748 METFORMIN ON AUTOPHAGY AND PERK. *Mol Carcinog* 2016;**55**:346–56.
749 doi:10.1002/mc.22284
- 750 99 Zhao C, Yin S, Dong Y, *et al.* Autophagy-dependent EIF2AK3 activation compromises
751 ursolic acid-induced apoptosis through upregulation of MCL1 in MCF-7 human breast
752 cancer cells. *Autophagy* 2013;**9**:196–207. doi:10.4161/auto.22805
- 753 100 Laplante M, Sabatini DM. mTOR Signaling in Growth Control and Disease. *Cell*
754 2012;**149**:274–93. doi:10.1016/j.cell.2012.03.017
- 755 101 B'chir W, Maurin A-C, Carraro V, *et al.* The eIF2 α /ATF4 pathway is essential for stress-
756 induced autophagy gene expression. *Nucleic Acids Res* 2013;**41**:7683–99.
757 doi:10.1093/nar/gkt563
- 758 102 Avivar-Valderas A, Salas E, Bobrovnikova-Marjon E, *et al.* PERK Integrates Autophagy and
759 Oxidative Stress Responses To Promote Survival during Extracellular Matrix Detachment.
760 *Mol Cell Biol* 2011;**31**:3616–29. doi:10.1128/MCB.05164-11
- 761 103 Rouschop KMA, van den Beucken T, Dubois L, *et al.* The unfolded protein response
762 protects human tumor cells during hypoxia through regulation of the autophagy genes
763 MAP1LC3B and ATG5. *J Clin Invest* 2010;**120**:127–41. doi:10.1172/JCI40027
- 764 104 Rzymiski T, Milani M, Pike L, *et al.* Regulation of autophagy by ATF4 in response to severe
765 hypoxia. *Oncogene* 2010;**29**:4424–35. doi:10.1038/onc.2010.191

- 766 105 Gade P, Manjegowda SB, Nallar SC, *et al.* Regulation of the Death-Associated Protein
767 Kinase 1 Expression and Autophagy via ATF6 Requires Apoptosis Signal-Regulating Kinase
768 1. *Mol Cell Biol* 2014;**34**:4033–48. doi:10.1128/MCB.00397-14
- 769 106 Gade P, Ramachandran G, Maachani UB, *et al.* An IFN- γ -stimulated ATF6-C/EBP- β -
770 signaling pathway critical for the expression of Death Associated Protein Kinase 1 and
771 induction of autophagy. *Proc Natl Acad Sci U S A* 2012;**109**:10316–21.
772 doi:10.1073/pnas.1119273109
- 773 107 Lopes F, Keita ÅV, Saxena A, *et al.* ER-stress mobilization of death-associated protein
774 kinase-1-dependent xenophagy counteracts mitochondria stress-induced epithelial
775 barrier dysfunction. *J Biol Chem* 2018;**293**:3073–87. doi:10.1074/jbc.RA117.000809
- 776 108 Appenzeller-Herzog C, Hall MN. Bidirectional crosstalk between endoplasmic reticulum
777 stress and mTOR signaling. *Trends Cell Biol* 2012;**22**:274–82.
778 doi:10.1016/j.tcb.2012.02.006
- 779 109 Yamazaki H, Hiramatsu N, Hayakawa K, *et al.* Activation of the Akt-NF-kappaB pathway
780 by subtilase cytotoxin through the ATF6 branch of the unfolded protein response. *J*
781 *Immunol Baltim Md 1950* 2009;**183**:1480–7. doi:10.4049/jimmunol.0900017
- 782 110 Hart LS, Cunningham JT, Datta T, *et al.* ER stress-mediated autophagy promotes Myc-
783 dependent transformation and tumor growth. *J Clin Invest* 2012;**122**:4621–34.
784 doi:10.1172/JCI62973
- 785 111 Li J, Ni M, Lee B, *et al.* The unfolded protein response regulator GRP78/BiP is required
786 for endoplasmic reticulum integrity and stress-induced autophagy in mammalian cells.
787 *Cell Death Differ* 2008;**15**:1460–71. doi:10.1038/cdd.2008.81
- 788 112 Ogata M, Hino S, Saito A, *et al.* Autophagy is activated for cell survival after endoplasmic
789 reticulum stress. *Mol Cell Biol* 2006;**26**:9220–31. doi:10.1128/mcb.01453-06
- 790 113 Wang W, Kang H, Zhao Y, *et al.* Targeting autophagy sensitizes BRAF-mutant thyroid
791 cancer to vemurafenib. *J Clin Endocrinol Metab* 2016;**;**jc.2016-1999. doi:10.1210/jc.2016-
792 1999
- 793 114 Keestra-Gounder AM, Byndloss MX, Seyffert N, *et al.* NOD1 and NOD2 signalling links ER
794 stress with inflammation. *Nature* 2016;**532**:394–7. doi:10.1038/nature17631
- 795 115 Kaneko M, Niinuma Y, Nomura Y. Activation Signal of Nuclear Factor- κ B in Response to
796 Endoplasmic Reticulum Stress is Transduced via IRE1 and Tumor Necrosis Factor
797 Receptor-Associated Factor 2. *Biol Pharm Bull* 2003;**26**:931–5. doi:10.1248/bpb.26.931
- 798 116 Urano F, Wang X, Bertolotti A, *et al.* Coupling of Stress in the ER to Activation of JNK
799 Protein Kinases by Transmembrane Protein Kinase IRE1. *Science* 2000;**287**:664–6.
800 doi:10.1126/science.287.5453.664

- 801 117 Castillo K, Rojas-Rivera D, Lisbona F, *et al.* BAX inhibitor-1 regulates autophagy by
802 controlling the IRE1 α branch of the unfolded protein response. *EMBO J* 2011;**30**:4465–78.
803 doi:10.1038/emboj.2011.318
- 804 118 Ding W-X, Ni H-M, Gao W, *et al.* Linking of Autophagy to Ubiquitin-Proteasome System
805 Is Important for the Regulation of Endoplasmic Reticulum Stress and Cell Viability. *Am J*
806 *Pathol* 2007;**171**:513–24. doi:10.2353/ajpath.2007.070188
- 807 119 Moretti J, Roy S, Bozec D, *et al.* STING Senses Microbial Viability to Orchestrate Stress-
808 Mediated Autophagy of the Endoplasmic Reticulum. *Cell* Published Online First: October
809 2017. doi:10.1016/j.cell.2017.09.034
- 810 120 NHS CB. 2013/14 NHS Standard Contract for Colorectal: Complex Inflammatory Bowel
811 Disease (Adult). 2013.
- 812 121 Bassi A, Dodd S, Williamson P, *et al.* Cost of illness of inflammatory bowel disease in the
813 UK: a single centre retrospective study. *Gut* 2004;**53**:1471–8.
814 doi:10.1136/gut.2004.041616
- 815 122 Bernstein CN, Loftus EV Jr, Ng SC, *et al.* Hospitalisations and surgery in Crohn's disease.
816 *Gut* 2012;**61**:622–9. doi:10.1136/gutjnl-2011-301397
- 817 123 Matsuda C, Ito T, Song J, *et al.* Therapeutic effect of a new immunosuppressive agent,
818 everolimus, on interleukin-10 gene-deficient mice with colitis. *Clin Exp Immunol*
819 2007;**148**:348–59. doi:10.1111/j.1365-2249.2007.03345.x
- 820 124 Massey DC, Bredin F, Parkes M. Use of sirolimus (rapamycin) to treat refractory Crohn's
821 disease. *Gut* 2008;**57**:1294–6. doi:10.1136/gut.2008.157297
- 822 125 Dumortier J, Lapalus M-G, Guillaud O, *et al.* Everolimus for refractory Crohn's disease: A
823 case report: *Inflamm Bowel Dis* 2008;**14**:874–7. doi:10.1002/ibd.20395
- 824 126 Mutalib M, Borrelli O, Blackstock S, *et al.* The use of sirolimus (rapamycin) in the
825 management of refractory inflammatory bowel disease in children. *J Crohns Colitis*
826 2014;**8**:1730–4. doi:10.1016/j.crohns.2014.08.014
- 827 127 Reinisch W, Panés J, Lémann M, *et al.* A multicenter, randomized, double-blind trial of
828 everolimus versus azathioprine and placebo to maintain steroid-induced remission in
829 patients with moderate-to-severe active Crohn's disease. *Am J Gastroenterol*
830 2008;**103**:2284–92. doi:10.1111/j.1572-0241.2008.02024.x
- 831 128 Ha J, Kim J. Novel pharmacological modulators of autophagy: an updated patent review
832 (2012-2015). *Expert Opin Ther Pat* 2016;**26**:1273–89.
833 doi:10.1080/13543776.2016.1217996
- 834 129 Galluzzi L, Bravo-San Pedro JM, Levine B, *et al.* Pharmacological modulation of
835 autophagy: therapeutic potential and persisting obstacles. *Nat Rev Drug Discov*
836 2017;**16**:487–511. doi:10.1038/nrd.2017.22

- 837 130 Rauthan M, Pilon M. A chemical screen to identify inducers of the mitochondrial
838 unfolded protein response in *C. elegans*. *Worm* 2015;**4**.
839 doi:10.1080/21624054.2015.1096490
- 840 131 Boyce M, Bryant KF, Jousse C, *et al*. A selective inhibitor of eIF2alpha dephosphorylation
841 protects cells from ER stress. *Science* 2005;**307**:935–9. doi:10.1126/science.1101902
- 842 132 Okazaki T, Nishio A, Takeo M, *et al*. Inhibition of the dephosphorylation of eukaryotic
843 initiation factor 2 α ameliorates murine experimental colitis. *Digestion* 2014;**90**:167–78.
844 doi:10.1159/000366414
- 845 133 Crespo I, San-Miguel B, Prause C, *et al*. Glutamine treatment attenuates endoplasmic
846 reticulum stress and apoptosis in TNBS-induced colitis. *PLoS One* 2012;**7**:e50407.
847 doi:10.1371/journal.pone.0050407
- 848 134 Yang L, Sha H, Davisson RL, *et al*. Phenformin activates the unfolded protein response in
849 an AMP-activated protein kinase (AMPK)-dependent manner. *J Biol Chem*
850 2013;**288**:13631–8. doi:10.1074/jbc.M113.462762
- 851 135 Kim H, Moon SY, Kim J-S, *et al*. Activation of AMP-activated protein kinase inhibits ER
852 stress and renal fibrosis. *Am J Physiol Renal Physiol* 2015;**308**:F226–236.
853 doi:10.1152/ajprenal.00495.2014
- 854 136 Hwang H-J, Jung TW, Ryu JY, *et al*. Dipeptidyl peptidase-IV inhibitor (gemigliptin) inhibits
855 tunicamycin-induced endoplasmic reticulum stress, apoptosis and inflammation in H9c2
856 cardiomyocytes. *Mol Cell Endocrinol* 2014;**392**:1–7. doi:10.1016/j.mce.2014.04.017
- 857 137 Yusta B, Baggio LL, Estall JL, *et al*. GLP-1 receptor activation improves beta cell function
858 and survival following induction of endoplasmic reticulum stress. *Cell Metab* 2006;**4**:391–
859 406. doi:10.1016/j.cmet.2006.10.001
- 860 138 Engin F, Hotamisligil GS. Restoring endoplasmic reticulum function by chemical
861 chaperones: an emerging therapeutic approach for metabolic diseases. *Diabetes Obes*
862 *Metab* 2010;**12 Suppl 2**:108–15. doi:10.1111/j.1463-1326.2010.01282.x
- 863 139 Allaire JM, Crowley SM, Law HT, *et al*. The Intestinal Epithelium: Central Coordinator of
864 Mucosal Immunity. *Trends Immunol* 2018;**39**:677–96. doi:10.1016/j.it.2018.04.002
- 865 140 Boyapati RK, Rossi AG, Satsangi J, *et al*. Gut mucosal DAMPs in IBD: from mechanisms to
866 therapeutic implications. *Mucosal Immunol* 2016;**9**:567–82. doi:10.1038/mi.2016.14
- 867 141 Boyapati RK, Dorward DA, Tamborska A, *et al*. Mitochondrial DNA Is a Pro-Inflammatory
868 Damage-Associated Molecular Pattern Released During Active IBD. *Inflamm Bowel Dis*
869 Published Online First: 1 May 2018. doi:10.1093/ibd/izy095
- 870 142 Odenwald MA, Turner JR. The intestinal epithelial barrier: a therapeutic target? *Nat Rev*
871 *Gastroenterol Hepatol* 2017;**14**:9–21. doi:10.1038/nrgastro.2016.169

- 872 143 Nishino K, Nishida A, Inoue R, *et al.* Analysis of endoscopic brush samples identified
873 mucosa-associated dysbiosis in inflammatory bowel disease. *J Gastroenterol* 2018;**53**:95–
874 106. doi:10.1007/s00535-017-1384-4
- 875 144 Wildenberg ME, Koelink PJ, Diederens K, *et al.* The ATG16L1 risk allele associated with
876 Crohn's disease results in a Rac1-dependent defect in dendritic cell migration that is
877 corrected by thiopurines. *Mucosal Immunol* 2017;**10**:352–60. doi:10.1038/mi.2016.65
- 878 145 Wildenberg M, Levin A, Vos C, *et al.* P668 ATG16L1 genotype is associated with response
879 to anti-TNF in vitro. *J Crohns Colitis* 2013;**7**:S279. doi:10.1016/s1873-9946(13)60689-3
- 880 146 Levin AD, Koelink PJ, Bloemendaal FM, *et al.* Autophagy Contributes to the Induction of
881 Anti-TNF Induced Macrophages. *J Crohns Colitis* 2016;**10**:323–9. doi:10.1093/ecco-
882 jcc/jjv174
- 883 147 Vos ACW, Wildenberg ME, Arijns I, *et al.* Regulatory macrophages induced by infliximab
884 are involved in healing in vivo and in vitro. *Inflamm Bowel Dis* 2012;**18**:401–8.
885 doi:10.1002/ibd.21818
- 886 148 Niess JH, Klaus J, Stephani J, *et al.* NOD2 polymorphism predicts response to treatment
887 in Crohn's disease--first steps to a personalized therapy. *Dig Sci* 2012;**57**:879–86.
888 doi:10.1007/s10620-011-1977-3
- 889 149 Chouchana L, Fernández-Ramos AA, Dumont F, *et al.* Molecular insight into thiopurine
890 resistance: transcriptomic signature in lymphoblastoid cell lines. *Genome Med* 2015;**7**:37.
891 doi:10.1186/s13073-015-0150-6
- 892 150 Ventham NT, Kennedy NA, Nimmo ER, *et al.* Beyond gene discovery in inflammatory
893 bowel disease: the emerging role of epigenetics. *Gastroenterology* 2013;**145**:293–308.
894 doi:10.1053/j.gastro.2013.05.050
- 895 151 Kalla R, Ventham NT, Kennedy NA, *et al.* MicroRNAs: new players in IBD. *Gut*
896 2015;**64**:504–17. doi:10.1136/gutjnl-2014-307891
- 897 152 Stevens TW, Matheeuwsen M, Lönnkvist MH, *et al.* Systematic review: predictive
898 biomarkers of therapeutic response in inflammatory bowel disease--Personalised
899 medicine in its infancy. *Aliment Pharmacol Ther* Published Online First: 30 October 2018.
900 doi:10.1111/apt.15033
- 901 153 Pascal V, Pozuelo M, Borruel N, *et al.* A microbial signature for Crohn's disease. *Gut*
902 2017;;gutjnl-2016-313235. doi:10.1136/gutjnl-2016-313235

903